



# Toxaphene 40 Wettable

[INSECTICIDE]

NET WEIGHT 50 LBS.

## For Spray Applications

Active Ingredient	By Wt.
Toxaphene*	40%
Inert Ingredients	60%
*Technical Chlorinated Camphene	
Chlorine Content—67% to 69%	

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

TURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE.

### DIRECTIONS

Apply to crops within 30 days of harvest unless otherwise directed. Residues remaining at harvest should be removed by washing, wiping, or other effective means. ORTHO Toxaphene 40 Wettable has enhanced killing power through the use of a wettable concentrate with lead arsenate, DDT and Dieldrin.

**BEANS:** Grasshoppers, Crickets—4 lbs. per acre in water to cover. Make one application to Green or Snap Beans after the pods start to form. No further application is necessary. Do not apply to wet or recently treated areas. Do not apply to beans intended for seed.

**BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, CELERY:** Climbing Cutworms, Cabbage Caterpillars, Thrips, Serpentine Leaf Miner—2½ to 5 lbs. per acre in water to cover. Concentrate Sprayer—4 lbs. per acre in water to cover. Grasshoppers, Crickets—4 lbs. per acre in water to cover. Do not




DO NOT INHALE  
DO NOT GET ON SKIN  
DO NOT TAKE INTERNALLY

KEEP OUT OF REACH OF CHILDREN

**WARNING:** May be fatal if swallowed. Do not breathe dust or spray mist. Do not get in eyes, on skin or on clothing. Wash thoroughly after using. Keep away from children and domestic animals. Do not store or transport with feed or food. Do not use on young plants. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

Highly toxic to fish and wildlife. Birds feeding on treated areas may be killed. Do not contaminate lakes, ponds, streams, woodlands and other non-crop areas with this spray or washings from the spray tank. This product is toxic to bees and should not be applied when bees are actively visiting the treated area.



Apply to corn after tasseling and before silks appear. Make one application at tasseling and one after silks appear. Do not apply to corn intended for seed. Do not apply to corn intended for silage. Do not apply to corn intended for grain. Do not apply to corn intended for feed.

**CORN:** Corn Earworms—Treat at tasseling and after silks appear. Make one application at tasseling and one after silks appear. Budworms—1 to 2 lbs. per acre in water to cover. Grasshoppers, Crickets—4 lbs. per acre in water to cover. Do not feed treated forage to any animals or animals to be finished for slaughter.

**PEPPERS:** Grasshoppers, Crickets—4 lbs. per acre in water to cover. Serpentine Leaf Miner—2½ lbs. per acre in water to cover. Do not apply within 5 days of harvest.

**TOMATOES:** Tomato Fruitworm, Hornworm, Stink Bugs, Thrips, Leaf Miners, Climbing Cutworms—5 lbs. per acre in water to cover. Concentrate Sprayer—4 to 5 lbs. per acre in water to cover. Grasshoppers, Crickets—4 lbs. per acre in water to cover. Do not apply within 1 day of harvest.

**GRASSHOPPERS and CRICKETS on listed crops:** Apply at recommended dosages per acre, in sufficient water for even distribution. Treat roadsides, field margins, idle land, and the fields to be used when main batch is completed or when young hoppers begin to move.

**BEEF CATTLE, SHEEP, GOATS:** Ticks—2 lbs. per 100 gal. water. Treat young animals lightly and with caution. Do not apply to animals under 2 months of age. Do not spray or dip animals when weather is cold or stormy, as injury may result from chilling. Do not spray or dip thirsty, overheated or emaciated animals. Do not use on dairy animals or inside dairy barns. Do not use within 4 weeks of slaughter.

**BARN AND OTHER BUILDINGS:** Flies—20 lbs. per 100 gal. water. Wet cracks thoroughly where insects alight or crawl. Cover feed or food utensils and drinking troughs before treatment. Do not apply this dosage to animals. Do not apply in dairy barns, milk rooms or poultry houses. Repeat as necessary to give control.

**CONDITIONS OF SALE:** Chevron Chemical Company, Chevy Chase, Maryland, warrants that the material conforms to the chemical description on the label and is reasonably fit for the use and the brand shown on the label. This warranty is made by the manufacturer and represents the best knowledge and belief of the manufacturer at the time the material was produced.

This material is sold as a pesticide and is not to be used for any other purpose. It is not to be used in any manner that is prohibited by law. It is not to be used in any manner that is prohibited by the manufacturer. It is not to be used in any manner that is prohibited by the user. It is not to be used in any manner that is prohibited by the law. It is not to be used in any manner that is prohibited by the manufacturer. It is not to be used in any manner that is prohibited by the user. It is not to be used in any manner that is prohibited by the law.

## CHEVRON CHEMICAL COMPANY

ORTHO DIVISION — San Francisco, California 94120

FORM 1081 J2

USDA Reg. No. 239-211

Product No. 1091

Made in U.S.A.

RICHMOND, CALIFORNIA

PORTLAND, OREGON

FRESNO, CALIFORNIA

DES MOINES, IOWA

CHERRY HILL, NEW JERSEY

DALLAS, TEXAS

HONOLULU, HAWAII

ORLANDO, FLORIDA