

Toxaphene 10 Dust

[INSECTICIDE]

NET WEIGHT 50 LBS.

Active Ingredient
Toxaphene*
Inert Ingredients

*Technical Chlorinated Camphene
(Chlorine Content: 67% to 69%)

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER.

BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE.


DIRECTIONS

This product is formulated for application by ground or aerial equipment.

ALFALFA SEED CROPS: Lygus Bugs, Grasshoppers, Thrips, Yellow Striped Armyworms, Alfalfa Loopers, Alfalfa Webworms and Tortrix Moth. 25 to 30 lbs. per acre, which includes 10 lbs. per acre for storage or feed.

BEANS: Lygus Bugs. Do not apply to beans when they are in treatment stage. Corn Earworms. 20 to 30 lbs. per acre. Do not apply to Green Beans, Lima Beans, or Broad Beans. Do not feed treated beans to swine, poultry or other animals which are fed for slaughter.

BROCCOLI, CAULIFLOWER, CABBAGE, BRUSSELS SPROUTS: Cabbage Caterpillars, Cabbage Looper. 20 lbs. per acre. Do not apply to Broccoli, Brussels Sprouts and



READ THE LABEL

CAUTION: Harmful if Swallowed! Avoid breathing dust. Avoid contact with eyes, skin and clothing. Wash thoroughly after using. Do not store or transport near feed or food. Do not use on young plants. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption. Highly toxic to fish and wildlife. Birds feeding on treated areas may be killed. Do not contaminate lakes, ponds, streams, woodlands and other non crop areas. This product is toxic to bees and should not be applied when bees are actively visiting the area.

cauliflower after cabb. parts begin to form or within 30 days of harvest. Do not apply to Cabbage after heads start to form.

CELERY: Leafminer. 20 to 30 lbs. per acre. Apply when leaves of stalks show signs of damage. Do not apply after the plant is half grown or after it starts to bolt.

CORN: Armyworm, Flea Beetle. 20 to 30 lbs. per acre. Do not feed treated corn to dairy animals or other animals which are fed for slaughter.

COTTON: Rapid Plant Bug, Tarnished Plant Bug, Fleahoppers. 20 to 30 lbs. per acre. Do not apply to cotton which is being fed to swine, poultry or other animals which are fed for slaughter.

Yellow-striped Armyworms, Cutworms 30 to 35 lbs. per acre. Grasshoppers—25 to 30 lbs. per acre. Do not graze dairy animals or animals being finished for slaughter in fields treated late in the season.

LETTUCE: Loopers, Cutworms—Use 50 lbs. per acre. Do not apply after seedling stage on leaf lettuce. Do not apply on head lettuce after heads start to form.

PEPPERS: Hornworms, Cutworms, Loopers, Beet Armyworms 30 lbs. per acre. Do not apply within 5 days of harvest.

TOMATOES: Leaf Miner, Thrips, Armyworms 35 to 40 lbs. per acre. Do not apply within 3 days of harvest.

CONDITIONS OF SALE: 1. Chevron Chemical Company. Chevron warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed herein. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or MERCHANTABILITY, guaranteed or representation, express or implied, concerning this material.

2. Material and bags are supplied by Chevron Chemical Company. Do not use this material unless you have contacted with the use of chemical safety equipment, but you get credit for removal of equipment. 3. This material is supplied in 50 lb. bags and damage caused by moisture, other pests or insects. Such damage does not affect the product as represented. 4. For the use of this material, please refer to the label and instructions. 5. This material is not intended for use on crops which are fed to swine, poultry or other animals which are fed for slaughter.

Deleted Red Clover

CHEVRON CHEMICAL COMPANY

FORM 1042-12

USDA Reg. No. 239-241

ORTHO DIVISION — San Francisco, California 94120

Product No. 1088

Made in U.S.A.

RICHMOND, CALIFORNIA

PORTLAND, OREGON

FRESNO, CALIFORNIA



DES MOINES, IOWA

CHERRY HILL, NEW JERSEY

HONOLULU, HAWAII

ORLANDO, FLORIDA