READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

ORTHO Toxaphene 4.8 Emulsive is for agricultural and commercial pest control use.

DIRECTIONS

BROCCOLL CAULIFLOWER, CABBAGE, BRUSSELS SPROUTS: Cabbage Caterpillars, Cabbage Looper—11/4 ats. per acre in water to cover. Do not apply to Broccoli, Cauliflower and Brussels Sprouts after the edible parts start to form or within 30 days of harvest. Do not apply to cabbage after heads start to form.

GRASSHOPPERS AND CRICKETS on Corn, Lettuce, Sorahum, Soybeans and Tomatoes—Use 2½ to 3 pints of ORTHO Toxaphene 4.8 Emulsive per acre as a spray in sufficient water for even distribution. Treat road-sides, field margins, idle land and the fields themselves when main hatch is completed or when young hoppers begin to move. Treat margins of cornfields and adjacent land. Late in the season, when most of the grasshoppers are adults and vegetation is tall and dense, a slight increase in the dosage of both sprays and dust may be necessary. Do not feed treated Soybeans or Corn forage to dairy animals or animals being finished for slaughter. Do not apply to Sorghum within 28 days of har-

vest. Do not apply to Sorghum more than once after heads start to for Do not graze dairy animals or animals being finished for slaughted treated Sorghum fields and do not ensile treated forage. Do not a after seedling stage for leaf lettuce. Do not apply on head lettuce (heads start to form. Do not apply to Tomatoes within 1 day of har Do not apply to soybeans within 21 days of harvest. Do not feed tree soybean plants or ensilage made from treated plants to poultry, a animals or animals being finished for slaughter. Do not feed soyl mill trash to livestock or poultry.

ANTS—For control of common house-infesting Ants, including the Artine Ant, and Pharaoh's Ant or the Little Red Ant: Use 5 gallons is gallons of water to make 100 gallons of spray. Treat outside of build at ground level; along porches, walks, out-buildings; and near known ant nest and runways. Repeat at monthly intervals or as necessity.

FLIES—For Houseflies: Use 5 gallons in 95 gallons of water to make gallons of spray. Spray the outside of buildings, including porches, apply as a light mist spray without causing excessive run-off. Remonthly or as necessary.

WARNING: May Be Fatal If Swallowed! Do not breathe s mist. Do not get in eyes, on skin or on clothing. Wash thoroughly using. Keep away from children and domestic animals. Do not stor transport with feed or food. DO NOT USE, POUR, SPILL OR ST y to Sorghum more than once after heads start to form. liry animals or animals being finished for slaughter on fields and do not ensile treated forage. Do not apply uge for leaf lettuce. Do not apply on head lettuce after my Do not apply to Tomatoes within 1 day of harvest. Soybeans within 21 days of harvest. Do not feed treated or ensilage made from treated plants to poultry, dairy als being finished for slaughter. Do not feed soybean rock or poultry.

of common house-infesting Ants, including the Argenaraoh's Ant or the Little Red Ant: Use 5 gallons in 95 o make 100 gallons of spray. Treat outside of buildings along porches, walks, out-buildings; and near any and runways. Repeat at monthly intervals or as necessary.

lies: Use 5 gallons in 95 gallons of water to make 100 Spray the outside of buildings, including porches, etc., mist spray without causing excessive run-off. Repeat cessary.

IG: May Be Fatal If Swallowed! Do not breathe spray heyes, on skin or on clothing. Wash thoroughly after from children and domestic animals. Do not store or ed or food. DO NOT USE, POUR, SPILL OR STORE

NEAR HEAT OR OPEN FLAME. Do not use on young plants. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

Highly toxic to fish and wildlife: Birds feeding in treated areas may be killed. Do not contaminate any body of water or apply to areas or crops other than those for which directions are given on the label. This product is toxic to bees and should not be applied when bees are actively visiting the area.

CONDITIONS OF SALE: 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.


2. Critical and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Chevron under 1 above) resulting from handling, storage, and use of this material.

ints. Do not ige or other ndered unfit

reas may be to areas or the label en bees are

hevron) waron the label ither makes, her warranty entation, ex-

ntrol prevent micals. Such ind crops to and damage even though even though and assume



[INSECTICIDE]

Active Ingredients	By	Wt
Toxaphene*		50%
Petroleum Distillate		44%
Inert Ingredients		6%
*Technical Chlorinated Camphene		
(Chlorine Content 67% to 69%)		
Contains 4.8 lbs. Toxaphene per gal.		

DO NOT INHALE

DO NOT GET ON SKIN

DO NOT TAKE INTERNALLY

WARNING: KEEP OUT OF REACH OF CHILDREN. SEE BACK PANEL FOR ADDITIONAL CAUTIONS.

WHEN CONTAINER IS EMPTY, IMMEDIATELY WASH THOROUGHLY AND DESTROY. NEVER RE-USE.

NET CONTENTS 1 GALLON LIQUID