

ACCEPTED

OCT 22 1975

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 239-279



ORTHO VOLCK Paste Emulsion

Dilution Table

1 Tbs. = 1 Tablespoonful

Add Water to Make	1 to 50 (2%)	1 to 75 (1 1/3%)	1 to 100 (1%)
1 Gallon	5 1/4 Tbs.	3 1/2 Tbs.	3 Tbs.
3 Gallons	1/2 Pint	1/3 Pint	1/4 Pint
5 Gallons	5/6 Pint	1/2 Pint	2/5 Pint
100 Gallons	2 Gallons	1 1/2 Gallons	1 Gallon

VOLCK Paste Emulsion requires only dilution with water at required strength, with agitation, to be ready for spraying. Careful and thorough wetting is essential for effective control.

CAUTION: Harmful if swallowed. Wash hands after use.

Note to Physicians: Emergency Information—call (415) 233-3737. Never combine VOLCK with sulfur or lime-sulfur. Never use sulfur and VOLCK on the same plant within a 30-day period. Avoid spraying trees that are suffering from lack of moisture. Do not spray when temperature exceeds 90°. Do not use on Maiden-hair Ferns, Cocos Palms, Maples, etc. or non-woody plants unless specified. May cause discoloration to Blue Spruce. Do not re-use empty container. Destroy it by perforating or crushing. Bury or discard in a safe place away from water supplies.

Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

CODE 131

Chevron Chemical Company

Ortho Division/San Francisco CA 94119 Richmond CA 94804

Product 772 Made in U.S.A.

Form 2279-J

EPA Reg. No. 239-279

EPA Est. 239-FL-1 No 002263 JA



ORTHO VOLCK® Paste E

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH DIRECTIONS AND DIRECTIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONTAINER OR DILUTE INTO FOOD OR DRINK CONTAINERS.

DIRECTIONS: Do not store diluted spray. Make new dilution.

AZALEAS: As a spray for Red Spider Mites, Mealybugs, and Scale Insects, use 12 Tablespoonfuls (6 oz.) of VOLCK Paste Emulsion to 3 gallons of water (1 to 75). Make three treatments a year: (1) in spring just after bloom or early June, (3) late September.

CAMELIAS: For the spring application against Scale Insects and Mites, use 16 Tablespoonfuls (8 oz.) of VOLCK Paste Emulsion to 3 gallons of water (1 to 50). For the late September application use 12 Tablespoonfuls of VOLCK Paste Emulsion to 3 gallons of water (1 to 75).

CITRUS, AVOCADOS, MANGOES: As a spray for Scale Insects and Red Spider Mites. Use 12 Tablespoonfuls (6 oz.) of VOLCK Paste Emulsion to 3 gallons of water (1 to 75) at any time of year in accordance with label directions. This application is also effective for the removal of sooty mold and control of whitefly larvae.

For light infestations, the dosage may be reduced to 8 Tablespoonfuls of VOLCK Paste Emulsion to 3 gallons of water (1 to 100). For best results, use when immature or crawler stage is present.

VOLCK Paste Emulsion is also recommended for use against Scale Insects, Mealybugs, Red Spider Mites, on a number of the more hardy ornamental, fruit, nut, and evergreen trees and shrubs such as certain palms, ligustrums and pecans.

HIGHLY REFINED FOLIAGE OIL SPRAY

100 Gallons

2 Gallons

1 1/2 Gallons

1 Gallon

VOLCK Paste Emulsion requires only dilution with water at required strength, with agitation, to be ready for spraying. Careful and thorough wetting is essential for effective control.

CAUTION: Harmful if swallowed. Wash hands after use.

Note to Physicians: Emergency Information—call (415) 233-3737. Never combine VOLCK with sulfur or lime-sulfur. Never use sulfur and VOLCK on the same plant within a 30-day period. Avoid spraying trees that are suffering from lack of moisture. Do not spray when temperature exceeds 90°. Do not use on Maiden-hair Ferns, Cocos Palms, Maples, etc. or non-woody plants unless specified. May cause discoloration to Blue Spruce. Do not re-use empty container. Destroy it by perforating or crushing. Bury or discard in a safe place away from water supplies.

Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

CODE 131

Chevron Chemical Company

Ortho Division, San Francisco, CA 94119 Richmond, CA 94804

Product 772 Made in U.S.A.

Form 7779 J

EPA Est. 239 FL-1 No. 002263 IA

EPA Reg. No. 739-279

Emulsion

- 1 to 100 (1%)
- 3 Tbs
- 1/4 Pint
- 1/5 Pint
- 1 Gallon

Required strength. Wetting is essen

737 Never com-
 CK on the same
 ing from lack of
 use on Maiden-
 s specified May
 er Destroy it by
 water supplies
 aning of equip-
 on this label

It in accordance


02263 IA



ORTHO

VOLCK®

Paste Emulsion





ORTHO VOLCK Paste Emulsion *237-279*

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS AND DIRECTIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

DIRECTIONS: Do not store diluted spray. Make new dilution for each use.

AZALEAS: As a spray for Red Spider Mites, Mealybugs, and Scale Insects, use 12 Tablespoonfuls (6 oz.) of VOLCK Paste Emulsion to 3 gallons of water (1 to 75). Make three treatments a year: (1) in spring just after blooming, (2) late May or early June, (3) late September

CAMELIAS: For the spring application against Scale Insects and Red Spider Mites, use 16 Tablespoonfuls (8 oz.) of VOLCK Paste Emulsion to 3 gallons of water (1 to 50) For the late September application use 12 Tablespoonfuls (6 oz.) of VOLCK Paste Emulsion to 3 gallons of water (1 to 75).

CITRUS, AVOCADOS, MANGOES: As a spray for Scale Insects, Whiteflies and Red Spider Mites Use 12 Tablespoonfuls (6 oz.) of VOLCK Paste Emulsion to 3 gallons of water (1 to 75) at any time of year in accordance with local spray practice. This application is also effective for the removal of sooty mold fungus through the control of whitefly larvae


For light infestations, the dosage may be reduced to 8 Tablespoonfuls (4 oz.) of VOLCK Paste Emulsion to 3 gallons of water (1 to 100). For best control of scales, use when immature or crawler stage is present

VOLCK Paste Emulsion is also recommended for use against Scale Insects, Whiteflies, Red Spider Mites, on a number of the more hardy out-of-door grown ornamental, fruit, nut, and evergreen trees and shrubs such as gardenias, hibiscus, certain palms, figs, and pecans

HIGHLY REFINED FOLIAGE OIL SPRAY

chevron

ORTHO VOLCK Paste Emulsion



Active Ingredient	By Wt.
Petroleum Oil	80%
Inert Ingredients	20%
(Pet Oil — Min. U.R. 92%)	

CAUTION: Keep out of reach of children.
See back panel for additional cautions.

NET CONTENTS 5 GALLONS LIQUID

ste Emulsion


ORTHO VOLCK Paste Emulsion

ORTHO VOLCK

READ INSTRUCTIONS
FOR DIRECTION

FOR USE

FOR USE



Active Ingredient	By Wt
Petroleum Oil	80%
Inert Ingredients	20%
(Pet Oil - Mineral Oil)	92%

CAUTION: Keep out of reach of children.
See back panel for additional cautions.

NET CONTENTS 5 GALLONS LIQUID

ORTHO

VIOLIN

Ortho Violin



CAUTION Keep out of reach of children
See back panel for additional cautions.

NET CONTENTS 5 GALLONS (18.9 L)