


ZINC COPOSOIL® 20 Dust



(FUNGICIDE)

Active Ingredient	By Wt.
Copper (in basic cupric zinc sulfate complex) expressed as metallic	3.6%
Inert Ingredients (Including 3.6% Zinc as Metallic*)	96.4%
*Zinc derived from basic cupric zinc sulfate complex.	

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER

KEEP OFFICE AND HOME WHEN EMERGENCY STAY OUT OF SMOKE

DIRECTIONS

Apply before the disease is expected to appear, as determined by State Agricultural Experiment Stations or by previous experience in the locality concerned. Dust the plants thoroughly. This product is formulated for application by ground or air treatment.

NET WEIGHT 50 LBS.

Chevron Chemical Company

2000 D Street, Emeryville, Calif. 94119
 Palmdale, California Fresno, California Honolulu, Hawaii
 Des Moines, Iowa Perth Amboy, New Jersey Orlando, Florida

Form 208 W Rev. 1-68-71 Made in U.S.A.

EPA Reg. No. 2-30 AA

CELERY: Blight—30 to 40 lbs. per acre. Apply when blight appears. Repeat at 7 to 10-day intervals.

PEARS: Fire Blight (Pear Blight)—20 to 35 lbs. per acre. Begin dusting at 15 to 20% bloom and repeat at 5 to 7-day intervals during bloom. Especially important just prior to rainy periods and when rat-tail blossoms are present.

POTATOES: Late Blight—Use 25 to 50 pounds per acre depending on plant size. Dust thoroughly and repeat the applications at 7-day intervals, as necessary.

SUGAR BEETS: Cercospora Leaf Spot—50 to 60 lbs. per acre. Apply at first sign of disease and repeat as necessary.

TOMATOES: Late Blight—Use 25 to 35 pounds per acre depending on plant size. Dust thoroughly

before disease appears, usually starting before early fall rains are expected. Repeat at 7 to 10-day intervals.

WALNUTS: Walnut Blight—60 to 70 lbs. per acre. Apply during early or late pre-bloom. Make 2 to 4 applications, the first at early pre-bloom stage (1% of blossoms showing), the second application at pre-bloom stage (10 to 20% of blossoms showing). Also 1 or 2 post-bloom dustings as may be required during damp weather.

CONDITIONS OF SALE: Chevron Chemical Company Chevron warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Certain and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants or crops to which the material is applied, lack of compatibility, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and, though label directions are followed, the user uses and uses a knowledge and common sense and skill to avert those assumed by the user. Chevron is not liable for any damage or loss resulting from the use of this material.

CAUTION READ THE LABEL

KEEP OUT OF REACH OF CHILDREN

CAUTION: Harmful if swallowed! Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

This product is toxic to fish. Keep out of any body of water.