

ORTHO®

[INSECTICIDE]

NET CONTENTS LIQUID

Active Ingredients	By Wt.
Toxaphene*	71%
Petroleum Distillate	16%
Inert Ingredients	13%
*Technical Chlorinated Camphene (Chlorine Content)	67% to 69%

Contains 8 pounds Toxaphene per gallon.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

WASH AND DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

Net contents of this container are approximately:

USE DESIGNATED AMOUNT OF PRODUCT IN THE FOLLOWING GALLONS WATER PER ACRE UNLESS OTHERWISE NOTED:

GROUND APPLICATION—Vegetable and Field Crops—10 to 20 gal's
Cotton—2 to 20 gal's Deciduous Fruit Crops—60 to 100 gal's

Where Recommended for AIR APPLICATION—Field Crops—2 to 20 gal's
Orchard Crops—10 to 20 gal's

ANTS—For control of common house-infesting Ants, including the Argentine Ant, Pharaoh's Ant, Little Red Ant: Use 2½ gallons in sufficient water to make 100 gallons of spray. Treat outside of buildings at ground level; apply periodic weekly treatments around perimeter. Do not apply to food or water. Repeat at monthly intervals or as necessary.

BEANS (including dry, green and lima): Climbing Cutworms, Darkling Ground beetle, Corn Earworm, Lygus Bugs—1 to 2 quarts per acre in water to cover. May be applied to green or mature beans after the pods start to form. No limitation on use of treated beans as human food. Do not feed to livestock. Do not apply to beans intended for slaughter.

BROCCOLI, CAULIFLOWER, BRUSSELS SPROUTS: Cabbage Caterpillars, Cabbage Loopers—1 to 4 quarts per acre in water to cover. Do not apply to Broccoli, Cauliflower and Brussels Sprouts after the heads have started to form or within 30 days of harvest.

BROCCOLI, CAULIFLOWER: Cutworms (Climbing or Surface Feeding)—4 quarts per acre in water to cover. Do not apply after edible parts start to form or within 30 days of harvest.

CABBAGE: Cabbage Caterpillars, Cabbage Loopers—1 to 2 quarts per acre in water to cover. Do not apply after heads start to form.



DO NOT TAKE INTERNALLY
DO NOT INHALE
DO NOT GET ON SKIN

KEEP OUT OF REACH OF CHILDREN

WARNING:

May Be Fatal if Swallowed! Do not breathe spray mist. Do not get in eyes, on skin or on clothing. Wash thoroughly after using. Keep away from children and domestic animals. Do not store or transport near feed or food. DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME. Do not use on young plants. Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

Highly toxic to fish and wildlife. Do not contaminate lakes, ponds, streams, wetlands, and other non-crop areas. Do not contaminate water by cleaning of equipment or emptying containers. Do not apply where runoff is likely to occur. End Use: For use on agricultural crops. Do not use on crops intended for human consumption. Do not apply to crops intended for human consumption. Do not use on crops intended for human consumption.

CARROTS: Climbing Cutworms—1 quart per acre.

CELERY: Cutworms, Fall Armyworms, Thrips—2 quarts per acre. Do not apply after plants start to bunch or after plants are half mature.

COTTON: Climbing Cutworms, Armyworms, Thrips, Boll Weevil, Fleahoppers, Cotton Boll Worm, Leafworms, Lygus and Rapid Plant Bugs, Stink Bugs and Grasshoppers: Apply 1 quart per acre. Apply at 5-day intervals for Boll Weevil, and at 7 to 10 day intervals when Boll Weevil is not a factor in pest status. Do not graze dairy animals or animals being prepared for slaughter in fields treated later in the season. West—Cutworms, Yellowstriped Armyworms, Grasshoppers, Lygus Bug, Fleahopper, Thrips: 1 to 4 quarts per acre. Do not apply after plants are open. Use dilute solution in sufficient water for coverage.

FLIES: For Houseflies—Use 2½ gallons in sufficient water to make 100 gallons of spray. Spray the outside of buildings, including porches, and other areas where flies are a problem. Do not use on crops intended for human consumption.

GRASSHOPPERS AND CRICKETS on listed crops: Use 1½ pints per acre in spray or in water to cover. Do not apply to crops intended for human consumption. Do not apply to crops intended for human consumption. Do not apply to crops intended for human consumption.

LETTUCE: Cutworms, Cabbage Loopers, Armyworms—Use 3 to 5 pints per acre in water to cover. Do not apply after seedling stage on leaf lettuce. Do not apply on head lettuce after heads begin to form.

LYGUS BUGS, THRIPS, ARMYWORMS, ALFALFA LOOPERS, LEAFHOPPERS, CLOVER HEAD CATERPILLARS, WEBWORMS: Alfalfa (Seed)—3 to 3½ pints per acre. May be combined with 1 quart of Ortho Supreme Spray per acre in 20 to 30 gal's water by ground application or 10 to 15 gal's water by air. Do not feed treated alfalfa forage to dairy animals or animals being finished for slaughter.

ONION THRIPS: Onions—1 qt. per acre. Repeat at 7-day intervals for 2 more applications. Do not apply to green or spring onions.

PEACHES, PEARS, APPLES: Thrips, Pear Psylla, Lygus, Other Cat-Facing Insects, Oriental Fruit Moth—1 pint per 100 gal's water. Do not apply within 40 days of harvest or after ground cover spray.

PEPPERS: Climbing Cutworms—1½ quarts per acre in water to cover and do not apply within 5 days of harvest; or 3 quarts per acre in water to cover and do not apply after Peppers begin to form.

STRAWBERRIES: Crown Borer, Leaf Roller, Tarnished Plant Bug, Strawberry Weevil, Climbing Cutworm—2 to 3 pt. per acre in 50 to 200 gal's water by ground application; 10 to 20 gal's water by air. Do not apply after fruit start to form.

TOMATOES: Tomato Fruitworm, Hornworm, Stink Bugs, Thrips, Leaf Miners, Climbing Cutworms, Corn Earworms, Armyworms, Tomato Pinworms, Flea Beetles, Russet Mites—1 to 2½ quarts per acre in 100 to 200 gal's water. For air application, not less than 10 gal's water per acre. Do not apply within 30 days of harvest.

LIVESTOCK

Do not treat Beef Cattle, Goats, Sheep or Swine within 28 days of slaughter. Do not use on dairy animals or its dairy barns. Do not treat calves under 3 months of age.

SHEEP TICKS: Sheep—As a Dip—1¼ quarts in 100 gal's water. As a Spray—2¼ quarts in 100 gal's water, spraying from ground during warm weather, when weeds are only a few inches long.

LICE: Beef Cattle, Swine—As a Spray—2¼ quarts in 100 gal's water. Do not use on Dairy Cattle, Sheep, Goats. As a Dip—2 quarts in 100 gal's water. Do not use on Dairy Cattle.

HORN FLIES: Beef Cattle—As a Spray—2¼ quarts in 100 gal's water, applying a wet 2½ quarts per acre to animal. Do not use on Dairy Cattle.

TICKS: Beef Cattle, Swine—As a Spray—2¼ quarts in 100 gal's water. Beef Cattle—As a Dip—1¼ quarts in 100 gal's water. Do not dip in water for more than 10 minutes. Do not use on Dairy Cattle.

CONDITIONS OF SALE

Ortho and Ortho Supreme are registered trademarks of Ortho Chemical Corporation. This product is sold under license from Ortho Chemical Corporation. The label and its contents are subject to change without notice. The manufacturer makes no warranty, expressed or implied, for use of this product on any crop or animal other than those specifically mentioned on the label.

Ortho Chemical Corporation, 1000 North 10th Street, Des Moines, Iowa 50319. This product is sold under license from Ortho Chemical Corporation. The label and its contents are subject to change without notice. The manufacturer makes no warranty, expressed or implied, for use of this product on any crop or animal other than those specifically mentioned on the label.