

# Toxaphene 6 Emulsive



# ORTHO

## (INSECTICIDE)

Active Ingredients	By Wt.
*Toxaphene .....	58%
Petroleum Distillate .....	29%
Inert Ingredients .....	13%
*Technical Chlorinated Camphene (Chlorine Content .....	67% to 69%)
Contains 6 lbs. Toxaphene per gallon.	

## WARNING: KEEP OUT OF REACH OF CHILDREN.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

WASH AND DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

Not for use or storage in or around the home.

### DIRECTIONS

USE DESIGNATED AMOUNT OF PRODUCT IN THE FOLLOWING GALLONS WATER PER ACRE UNLESS OTHERWISE NOTED:  
GROUND APPLICATION—Vegetable and Field Crops—30 to 250 gals. Cotton—3 to 20 gals.  
Where Recommended for AIR APPLICATION—Field Crops—3 to 10 gals.

**BROCCOLI, CAULIFLOWER, CABBAGE, BRUSSELS SPROUTS:** Cabbage Caterpillars, Cabbage Looper—1 qt. per acre in sufficient water to cover. Do not apply to Broccoli, Brussels Sprouts and Cauliflower after edible parts begin to form or within 30 days of harvest. Do not apply to Cabbage after heads start to form.

## NET CONTENTS 5 GALS. LIQUID

**Chevron Chemical Company**

Ortho Division/San Francisco CA 94119

Product 1340

Form 8998-C

EPA Reg. No. 239 432

Made in U.S.A.

Richmond CA 94804

239-432

**CORN: Armyworms, Grasshoppers, Chinch Bugs**—Apply 1½ qts. per acre in sufficient water for coverage. Apply as first insects appear. Repeat as necessary. Do not feed treated forage to dairy animals or animals being finished for slaughter.

**CORN (Preplant or Preemergence Only): Armyworms, Cutworms**—Use 1½ to 2 qts. of ORTHO Toxaphene 6 Emulsive suitably diluted with water per acre on a broadcast basis. Do not use as a post-emergence treatment. Do not graze dairy animals on stover. Do not graze meat animals on stover within 4 weeks of slaughter. No limitation on the use of grain.

**COTTON: Cotton Boll Weevil, Cotton Bollworm, Armyworms, Fleahopper, Thrips, Leafworm, Grasshoppers**—1-2/3 qts. per acre in water to cover. (By Air: 3 to 5 gals. water per acre and By Ground Rig: 5 to 15 gals. water per acre.) **Early Application for Control of Thrips and Fleahoppers only:** Use 2/3 qt. in sufficient water to cover, depending on method of application. Do not graze dairy animals or animals being finished for slaughter in fields treated late in the season.



DO NOT TAKE INTERNALLY

DO NOT INHALE

DO NOT GET ON SKIN

**WARNING:** May Be Fatal If Swallowed! Do not breathe spray mist. Do not get in eyes, on skin or on clothing. Wash thoroughly after using. Keep away from children and domestic animals.

**Note to Physicians:** Emergency Information: call (415) 233-3737. Do not store or transport near feed or food. **DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.** Do not use on young plants. **Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.**

Highly toxic to Fish and Wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

**LETTUCE: Cutworms (Surface Feeding and Climbing)**—1 to 1½ qts. per acre in water to cover. Do not apply after seedling stage on Leaf Lettuce. Do not apply to Head Lettuce after heads begin to form.

**PEPPERS: Cutworms**—2 qts. per acre in water to cover. Do not apply within 5 days of harvest.

### LIVESTOCK

Do not treat Beef Cattle, Goats, Sheep or Swine within 28 days of slaughter. Do not use on dairy animals or in dairy barns. Do not treat calves under 3 months of age.

**SHEEP TICKS: Sheep—As a Dip**—1½ qts. in 100 gals. water. **As a Spray**—3 qts. in 100 gals. water, spraying thoroughly during warm weather, when wool is only a few inches long.

**LICE: Beef Cattle, Swine—As a Spray**—3 qts. in 100 gals. water. Do not use on Dairy Cattle. **Sheep, Goats—As a Dip**—1 qt. in 100 gals. water. Do not use on Dairy Goats.

**HORN FLIES: Beef Cattle—As a Spray**—3 qts. in 100 gals. water, applying about 2 qts. per mature animal. Do not use on Dairy Cattle.

**TICKS: Beef Cattle, Swine—As a Spray**—3 qts. in 100 gals. water. **Beef Cattle—As a Dip**—1½ qts. in 100 gals. water. Do not dip heavy, fat cattle in hot weather. Do not use on Dairy Cattle.

**CABLE TYPE BACK RUBBERS FOR Range Cattle to Control Horn Flies, Mosquitoes**—2/3 pt. ORTHO Toxaphene 6 Emulsive to 1 gal. No. 2 fuel oil or furnace oil. Do not use on dairy animals. Do not treat Beef Cattle within 28 days of slaughter.

**CONDITIONS OF SALE:** 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material. 2. Critical and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants and crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Chevron under 1. above) resulting from handling, storage and use of this material.