


# ZINC COPOSIK<sup>®</sup> Sulfur 35-30 Dust



# ORTHO

## (FUNGICIDE)

Active Ingredients	By Wt.
Copper (in basic cupric zinc sulfate complex expressed as metallic) .....	6.3%
Sulfur .....	30.0%
Inert Ingredients (Including 6.3% Zinc as Metallic*) .....	63.7%
*Zinc derived from basic cupric zinc sulfate complex.	

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE.

### DIRECTIONS

In the use of this dust, it is advisable that applications be started before the disease is expected to appear, as determined by State Agricultural Experiment Stations or by previous experience in the locality concerned. Dust the plants thoroughly and repeat the applications as necessary. Use at the rate of 40 to 50 pounds per acre unless otherwise specified. This product is formulated for application by ground or air equipment.

**BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER:** Downy Mildew—25 to 50 lbs. per acre. Apply at first sign of disease and repeat at

7 to 10-day intervals to maintain control.

**CARROTS: Late Blight, Early Blight**—Apply when blight appears. Repeat at 7 to 10-day intervals.

**CELERY: Celery Blight**—Apply when blight appears. Repeat at 7 to 10-day intervals.

**CUCURBITS: Early and Late Blight, Downy Mildew, Powdery Mildew, Leaf Spot, Anthracnose**—30 to 45 lbs. per acre. Apply before diseases appear. Repeat at 7 to 10-day intervals.

**LETTUCE: Downy and Powdery Mildew**—30 to 35 lbs. per acre. Apply prior to or at first sign of disease. Repeat at regular 5 to 10-day intervals to maintain control.

**ONIONS: Downy Mildew**—35 to 50 pounds per acre. Apply before disease appears. Repeat at 7 to 10 day intervals.

**POTATOES: Late Blight**—Use 40 to 60 lbs. per acre. Repeat the applications at 7-day intervals, as necessary.

**SPINACH: Downy Mildew**—Apply prior to or at first sign of disease. Repeat at regular 5 to 10-day intervals to maintain control.

**TABLE GRAPES: Bunch Rot (Botrytis), Powdery Mildew**—Dust just after thinning, 25 pounds per acre, and a second dusting just previous to ripening.

**TOMATOES: Late Blight**—Use 40 to 60 lbs. per acre. Dust thoroughly before disease appears, usually starting before early fall rains are expected. Repeat at 7 to 10-day intervals. Because of the sulfur content of this dust, first obtain permission from your canner or processor before using it on tomatoes intended for sale to such canner or processor.

**CONDITIONS OF SALE:** 1. Chevron Chemical Company Chevron warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risk and liability, except those growing directly out of the fact of sale, for the use of this material on crops to which it is applied.

**CAUTION**      **READ THE LABEL**

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION:** Harmful if swallowed. Some fruits and other plants are susceptible to injury from sulfur under certain climatic conditions. The user is advised not to use sulfur on any crop unless local use has proved that sulfur does not damage crops in that locality. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

## NET WEIGHT 50 LBS.

### Chevron Chemical Company

Ortho Division, San Francisco, Calif. 94111  
 Richmond, California      Dallas, Texas      New York, New York  
 Des Moines, Iowa      Fort Worth, Texas      St. Louis, Missouri  
 St. Paul, Minnesota      Phoenix, Arizona      Portland, Oregon  
 Berkeley, California      Los Angeles, California      San Diego, California  
 El Paso, Texas      Houston, Texas      Memphis, Tennessee