



**POTATOES:** Colorado Potato Beetle, Flea Beetle, Leafhopper, Serpentine Leaf Miner, Grasshoppers, White Fly, Aphids, Potato Psyllid— $\frac{3}{4}$  pint per 100 gallons water—200 diluted gals. per acre. By Airplane— $\frac{3}{4}$  pint per acre in water to cover.

**PLUMS, PRUNES:** Mosby Plum Aphids, Red Spider Mites, Fruit Tree Leaf Rollers, Leafhoppers, Tussock Moth— $\frac{1}{2}$  pt. to 100 gals. water. San Jose Scale—In Dormant—1 pint (not over 5 pts. per acre) in combination with 3 to 4 gals. Dormant oil or  $1\frac{1}{2}$  gals. Superior type oil.

**SPINACH:** Aphids, Serpentine Leaf Miner— $\frac{1}{2}$  to 1 pint. By Air— $\frac{3}{4}$  pint per acre in water to cover.

**STRAWBERRY:** Leaf Roller, Red Spider Mite— $\frac{1}{2}$  pint. Repeat at 7 to 10-day intervals. By Air— $\frac{3}{4}$  pint per acre in water to cover.

**TOBACCO (Field):** Aphids and Suckfly— $\frac{1}{3}$  pint. Make no applications within 5 days of priming or 15 days of cutting. Avoid getting wet with plant juices when cutting. By Air— $\frac{3}{4}$  pint per acre in water to cover. (Seedbed): Aphids— $\frac{1}{2}$  pint.

**TOMATOES:** Flea Beetle, Leafhopper, Serpentine Leaf Miner, Grasshoppers, White Fly, Aphids— $\frac{1}{2}$  pint with regular fungicide program, using 200 gallons diluted spray per acre. By Air—1 pint per acre in water to cover.

**VETCH, ALFALFA, CLOVER, WHEAT, OATS, BARLEY:** Greenbug (Aphids), Red Spider, Armyworms, Grasshoppers— $\frac{1}{2}$  pint per acre in water to cover. Alfalfa Weevil Larvae— $\frac{3}{4}$  pint per acre. Apply when alfalfa is 12 to 14 inches high. For severe infestation use 1 pint and repeat application to stubble after first cutting.

**WALNUTS:** Frosted Scale, Calico Scale and Other Soft Scales— $\frac{1}{2}$  pint plus 2 gals. ORTHOL-K Flowable Medium in full dormant. Aphids, Red Spider Mites— $\frac{1}{2}$  pint as needed during summer. Oyster Shell Scale— $\frac{1}{3}$  pint in May or June. Do not apply after outer husks open.

**MOSQUITO LARVAE CONTROL** by Experienced Personnel of Mosquito Abatement Districts: 50 cubic centimeters ORTHOPHOS 4 Emulsive in 4 gals. water per acre, applied to breeding areas.

**AIRPLANE AND OTHER CONCENTRATED SPRAY USE**—Use a very fine mist spray and never apply as coarse droplet spray to avoid spot injury to plants and fruit.

**DOSAGES**—Avoid use of higher dosages than shown, since excessive dosages may mark fruit or foliage and especially smooth skin varieties such as prunes, plums, cherries and olives.

**NOTICE:** The statements made on this label, or by any of our agents concerning this material, are given for information only. They are believed to be true and accurate, but because conditions of use which are of critical importance are beyond its control, California Chemical Company does not make, nor does it authorize any agent or representative to make, any warranty, guarantee or representation, express or implied, concerning this material, except that it conforms to the chemical description on the label. Neither California Chemical Company nor the seller shall be held responsible in any manner for any personal injury or property damage or other type of loss resulting to the buyer to any person from the handling, storage or use of this material whether or not in accordance with directions. The buyer assumes all risk and liability resulting from such handling, storage or use and accepts and uses this material on these conditions.

239-547