

2 of 3

HASTE COPY OF LAST PRINTED BAG. (This replaces mat usually sent.)

ACCEPTED
MAR 16, 1970
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER FEDERAL LAW SUBJECT
TO ATTACHED COMMENTS.

239-612

ORTHO

DIRECTIONS

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

AGRICULTURAL CROPS


Use 2 to 3 lbs. per 100 gals. water (5 to 8 teaspoonfuls per 1 gallon water). Thoroughly cover all foliage and fruit. Do not apply more than the following lbs. of actual Methoxychlor per acre per application: Small fruits (except Strawberries) 7 lbs.; Garden Crops (except as noted) and Strawberries 2 1/4 lbs.; Field and forage crops 7 lbs. Start applications at first sign of infestation (for fruits, begin at petal-fall) and repeat at 7 to 14-day intervals as needed. Do not apply to any crop later than 7 days before harvest or grazing unless otherwise noted.

Ortho Methoxychlor 50 W/W in 1 lb. can

Cameraman's Note
Poor Copy

[INSECTICIDE]

ACTIVE INGREDIENT By Wt.
Methoxychlor, Technical.....50%
INERT INGREDIENTS.....50%
Contains 44% 2,2-bis (p-methoxyphenyl)-1,1,1-trichloroethane and 6% of other isomers and related compounds.




READ THE LABEL
KEEP OUT OF REACH OF CHILDREN

USE ON: APPLES, QUINCES, PEARS—Apple Maggot, Cankerworms, Codling Moth, Japanese Beetle, Leafhoppers, Pear

[INSECTICIDE]

ACTIVE INGREDIENT By Wt.
 Methoxychlor, Technical.....50%
INERT INGREDIENTS.....50%
 Total content to 44% 2,2-bis (p-methoxyphenyl)-1,1,1-trichloroethane and 6% of other isomers and related compounds.



READ THE LABEL

KEEP OUT OF REACH OF CHILDREN

CAUTION: Wash thoroughly after handling. ~~Avoid contamination of food and feedstuffs.~~ Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.
 Highly toxic to fish. Do not contaminate any body of water, by direct application, cleaning of equipment or disposal of wastes and containers.

DIRECTIONS ON SIDE PANEL

BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE.

NET WEIGHT 4 LBS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

AGRICULTURAL CROPS

Use 2 to 3 lbs. per 100 gals. water (5 to 8 teaspoonfuls per 1 gallon water). Thoroughly cover all foliage and fruit. Do not apply more than the following lbs. of actual Methoxychlor per acre per application: Small fruits (except Strawberries) 2 lbs.; Garden Crops (except as noted) and Strawberries 2 1/4 lbs.; Field and forage crops 2 lbs. Start applications at first sign of infestation (for fruits, begin at petal-fall) and repeat at 7 to 14-day intervals as needed. Do not apply to any crop later than 7 days before harvest or grazing unless otherwise noted.

FRUIT TREES: APPLES, QUINCES, PEARS—Apple Maggot, Cankerworms, Codling Moth, Japanese Beetle, Leafhoppers, Pear Slug, Tent Caterpillar. CHERRIES—Cankerworms, Cherry Fruit Fly, Cherry Fruitworm, Japanese Beetle, Leafhoppers, Pear Slug, Plum Curculio, Tent Caterpillar. APRICOTS, NECTARINES, PEACHES, PLUMS, PRUNES—Cankerworms, Japanese Beetle, Leafhoppers, Pear Slug, Plum Curculio, Tent Caterpillar. Do not apply to Apples, Pears, Quinces, Cherries, Plums and Prunes within 7 days of harvest and to Apricots, Nectarines and Peaches within 21 days of harvest.

SMALL FRUITS: BLUEBERRIES, CURRANTS—Japanese Beetle, Leafhoppers. CRANBERRIES—Cranberry Fruitworm, Japanese Beetle. GOOSEBERRIES, LOGANBERRIES—Japanese Beetle. GRAPE—Grape Berry Moth,

CONTINUED ON BACK PANEL

Cameraman's Note

Poor Copy

C. C. C. Co. M. I. Co.

Gooseberries and Grapes within 14 days of harvest. Do not apply to Blackberries, Loganberries, Raspberries, and Strawberries within 14 days of harvest; however, application may be made up to 8 days of harvest at a rate of 3 1/2 lbs. of Cerio per acre. *3 1/2 lbs Cerio*

GARDEN CROPS: ASPARAGUS—Asparagus Beetle, Fall Armyworm, Flea Beetle, Japanese Beetle. **BEANS**—BEANHORNED BEAN, BEAN—Bean Leaf Beetle, Fall Armyworm, Flea Beetle, Japanese Beetle, Leafhopper, Green Bean Beetle, Flea Beetle. **PEAS**—Pea Leafhopper, Flea Beetle, Japanese Beetle, Leafhopper, Green Bean Beetle, Flea Beetle. **PEPPERS**—Pepper Weevil, Flea Beetle, Japanese Beetle, Leafhopper, Green Bean Beetle, Flea Beetle. **PUMPKINS**—Pumpkin Beetle, Flea Beetle, Japanese Beetle, Leafhopper, Green Bean Beetle, Flea Beetle. **SPINACH**—Spinach Beetle, Flea Beetle, Japanese Beetle, Leafhopper, Green Bean Beetle, Flea Beetle. **TOMATOES**—Tomato Pinworm, Flea Beetle, Japanese Beetle, Leafhopper, Green Bean Beetle, Flea Beetle. **CUCUMBERS**—Cucumber Beetle, Flea Beetle, Japanese Beetle, Leafhopper, Green Bean Beetle, Flea Beetle. **EGGPLANT**—Eggplant Beetle, Flea Beetle, Japanese Beetle, Leafhopper, Green Bean Beetle, Flea Beetle. **PEPPERS**—Pepper Weevil, Flea Beetle, Japanese Beetle, Leafhopper, Green Bean Beetle, Flea Beetle. **POTATOES**—Colorado Potato Beetle, Flea Beetle, Japanese Beetle, Leafhopper. Do not apply to Calliflower, Corn, Potatoes, Peas, Kale and Rutabagas within 7 days of harvest. Do not apply to Beets, Carrots, Spinach, Squash, Carrots, Collards, Kale, Lettuce, Spinach and Turnips with 7 days of harvest. May be applied to Asparagus after 8 days before harvest. Move plants by pulling or bending. May be applied to Beans and Black-eyed Peas up to 8 days of harvest. Peas are not used for feed or forage. Do not apply to Calliflower within 8 days of harvest. Application may be made to Cantaloupe, Cucumbers, Eggplant, Kohlrabi, Peppers, Pumpkins, Squash, Summer Squash and Tomatoes up to 24 hours of harvest at a rate of 3 1/2 lbs. of Cerio per acre. May be used up to harvest on Potatoes. *3 1/2 CERIO*

FIELD AND FORAGE CROPS—Apply one pint of Cerio per acre. **BARLEY**—Barley Yellowfly, Flea Beetle, Japanese Beetle, Leafhopper, Green Bean Beetle, Flea Beetle. **BEANS**—Bean Leaf Beetle, Flea Beetle, Japanese Beetle, Leafhopper, Green Bean Beetle, Flea Beetle. **PEAS**—Pea Leafhopper, Flea Beetle, Japanese Beetle, Leafhopper, Green Bean Beetle, Flea Beetle. **CUCUMBERS**—Cucumber Beetle, Flea Beetle, Japanese Beetle, Leafhopper, Green Bean Beetle, Flea Beetle. **EGGPLANT**—Eggplant Beetle, Flea Beetle, Japanese Beetle, Leafhopper, Green Bean Beetle, Flea Beetle. **PEPPERS**—Pepper Weevil, Flea Beetle, Japanese Beetle, Leafhopper, Green Bean Beetle, Flea Beetle. **POTATOES**—Colorado Potato Beetle, Flea Beetle, Japanese Beetle, Leafhopper. Do not apply to Calliflower, Corn, Potatoes, Peas, Kale and Rutabagas within 7 days of harvest or grazing. *50 lb. Cerio*

GRAIN STORAGE BINS

GRAIN STORAGE BINS: Cecidid, Flat Grain Beetle, Saw-Toothed Grain Beetle, Lesser Grain Borer, Red Flour Beetle, Long-headed Flour Beetle, White Weevil, Granary Weevil, Confused Flour Beetle, Foreign Grain Beetle, Flour Beetles—To prevent infestation in empty storage bins, use Cerio in the grain. Apply 1 quart of 1 gallon dilute spray per 200 sq. ft. Treat before products are stored.

LIVESTOCK—FARM BUILDINGS

LIVESTOCK: Apply one pint of Cerio per acre. Rub in lightly to carry powder beneath the hair. Application to sheep, following shearing, gives temporary relief from sheep ticks (beds). All the above treatments are usually effective for about 3 weeks.

Application to sheep, following shearing, gives temporary relief from sheep ticks (beds). All the above treatments are usually effective for about 3 weeks.

FARM BUILDINGS: Stable Flies, Houseflies, Ticks—Use 1 lb. in 2 1/2 gal. water. Apply 1 gallon mixture to approx. 1000 sq. ft. of surface—treating walls, floors, cracks, crevices, baseboards, pet bedding, stalls and pens. Such deposits are usually effective for 3 to 5 weeks, unless removed by weathering or other means. Care should be taken to avoid contamination of milk handling equipment, feedstuffs, feed troughs and watering receptacles. Dairy animals should not be present while spraying. Do not use in poultry houses. This use should be supplemented with proper sanitary practices including spraying or removal of breeding or nesting areas near use in and around buildings.

CONDITIONS OF SALE: 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit for the intended use. Chevron neither warrants nor represents any agent or representative of Chevron, other warranty of FIT-NESS, MERCHANTABILITY, GUARANTEE or otherwise, express or implied, concerning this material. 2. Control and unforeseeable factors beyond Chevron's control prevent it from eliminating insects in connection with the use of certain products. Such factors include, but are not limited to, damage to plants and animals, which the material is applied to, and the control, and damage caused by other plants or animals. Such factors occur even though the product is reasonably fit for the uses stated on the label and although such factors are not intended by user and are not intended by Chevron under the best of circumstances. Chevron does not warrant, represent or make any claim of fitness for use of this material.

*Cameraman's Note
Poor Copy*

3(043)

COPY A

Do not store or transport near feed or food.

COPY B

Do not repeat this treatment at intervals of less than 3 weeks.