

[239-646]

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS. WASH AND DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.

Contains Premium Grade Malathion

ON FRUIT AND NUT CROPS AND ORNAMENTALS, APPLY THE SPECIFIED AMOUNT IN 100 GALS. OF WATER UNLESS OTHERWISE NOTED.

USE DESIGNATED AMOUNT OF PRODUCT IN THE FOLLOWING GALLONS WATER PER ACRE UNLESS OTHERWISE NOTED:

GROUND APPLICATION—Vegetable and Field Crops—30 to 250 gals. Cotton—3 to 20 gals. Deciduous Fruit Crops—60 to 800 gals. Citrus—400 to 3,000 gals. (On Young Fruit Plantings—Use a minimum of 15 gals. water per acre).

Where Recommended for AIR APPLICATION—Field Crops—3 to 10 gals. Orchard Crops—10 to 20 gals.

Apply Thorough, Full-Coverage Spray. Repeat as necessary unless otherwise specified. In Orchards—Codling Moth, Oriental Fruit Moth and Plum Curculio—Apply at petal fall and every 10 to 14 days thereafter until controlled.

ALFALFA: Grasshoppers, Aphids (including Spotted Alfalfa Aphid), Lygus Bugs, Leafhoppers, Spider Mites, Alfalfa Weevil Larvae, Spittlebugs; Vetch Bruchids on Alfalfa—1½ to 2 pts. per acre in water to cover. Armyworms—2 pts. per acre in water to cover. Apply to alfalfa in bloom only in the evening or early morning when bees are not working in the field or are not hanging on the outside of hives. May be grazed or harvested on day of application.

ALFALFA (Seed): Lygus Bugs, Leafhoppers, Spider Mites, Aphids—1 to 2 pints per acre in water to cover. Do not use during periods of bee activity. Combine with 1 quart VOLCK Supreme Spray per acre in 20 to 30 gals. water by ground application or 10 to 15 gals. water by air.

APPLES: Codling Moth, Plum Curculio, Red-Banded Leaf Roller—2 pts. Bud Moth, Forbes Scale, Woolly Aphid—1 pt. Tent Caterpillars (Eastern)—1 to 1½ pts. Mealybug, Spider Mites—1 to 2 pts. Green Apple Aphid, Rosy Apple Aphid—1½ pts. Malathion may cause fruit injury to McIntosh and Cortland varieties if spray is applied within 4 weeks of harvest. Do not apply within 3 days of harvest.

ASPARAGUS: Thrips—1½ to 2 pts. in water to cover 1 acre. Asparagus Beetles—2 pts. per acre in water to cover. Do not apply within 1 day of harvest.

BEANS: Spider Mites, Lygus, Mexican Bean Beetle, Japanese Beetle—1½ to 2¾ pts. per acre. Aphids, Cucumber Beetles, Leafhoppers—2 pts. per acre. Do not apply within 1 day of harvest.

BEETS: Aphids—1½ to 2 pts. per acre. Do not apply within 7 days of harvest.

BLUEBERRIES: Japanese Beetles—1½ pints in water to cover 1 acre. Apply when first berries turn blue and at 10-day intervals as long as necessary. Do not apply within 1 day of harvest.

BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, CABBAGE, KALE, MUSTARD GREENS, TURNIPS: Aphids, Imported Cabbage-worm, Cabbage Looper, Flea Beetles, Diamond-back Caterpillars—2 pts. per acre, making full coverage application every 7 days. Do not apply to Broccoli and Turnips within 3 days of harvest and to Brussels Sprouts, Cauliflower, Cabbage, Kale and Mustard Greens within 7 days of harvest.

CARROTS: Leafhoppers—2½ pts. per acre in water to cover. Do not apply within 7 days of harvest.

CELERY, LETTUCE, SPINACH: Aphids, Spider Mites, Leafhoppers—2 pts. per acre. Loopers (on lettuce)—3 pts. per acre. Do not apply to Head Lettuce, Celery and Spinach within 7 days of harvest and to Leaf Lettuce within 14 days of harvest.

CHERRIES: Black Cherry Aphid, Leaf Roller, Fruit Fly—1½ pts. Injury may occur on certain varieties of sweet cherries, particularly in Northwest. Do not apply within 3 days of harvest.

CITRUS (Grapefruit, Lemons, Limes, Oranges, Tangerines): California Red Scale, Yellow Scale, Purple Scale, Black Scale (single-brooded), Soft Scale, Citricola Scale—1 to 1½ pts. Florida Red Scale, Purple Scale—Use 1 qt. per 100 gals. water. Thrips—1¼ pts. per 100 gals. water per acre. Do not apply when trees are in bloom. Do not apply within 7 days of harvest. Do not use more than 25 lbs. actual Malathion per acre on citrus.

CLOVER: Aphids, Mites, Leafhoppers, Grasshoppers—1½ to 2 pts. per acre in water to cover. Do not apply to clover in bloom. May be grazed or harvested on day of application.

COTTON: Certain Spider Mites, Aphids, Leafhoppers, Whiteflies, Brown Cotton Leafworm, Cotton Leafworm, Cotton Leaf Perforator, Thrips—½ to 2 pts. in water to cover 1 acre. Boll Weevil, Lygus Bug—1 to 2 qts. in water to cover 1 acre. Repeat as necessary. May be applied up to harvest.

CRANBERRIES: Leafhoppers, Spittlebug Nymphs, Black-headed Fireworms, Cranberry Fruitworm—1½ pints per acre. Do not apply within 3 days of harvest.

CUCUMBERS, SQUASH, MELONS, PUMPKINS: Aphids, Spider Mites, Pickleworms—1½ pts. per acre. Do not apply to cucurbits unless plants are dry. Squash Vine Borers—1½ qts. per acre, applying weekly to stems and vines at base of plants. Do not apply to Cucumbers, Squash and Melons within 1 day of harvest and to Pumpkins within 3 days of harvest.

EGGPLANT: Spider Mites, Aphids—1 pt. Lace Bugs—1½ qts. per acre. Do not apply within 3 days of harvest.

FIGS: Dried Fruit Beetles, Vinegar Flies—Apply 2 quarts plus 1 to 2 gals. unsulfured Molasses per acre in water to cover. Do not apply within 3 days of harvest.

FILBERTS: Filbert Aphid, Apple Mealybug, Scale (crawler stage), Tingids, Eye-Spotted Bud Moth—1 pt. May be used up to harvest.

GRAPES: Leafhoppers, Spider Mites, Mealybugs—1½ pts. Injury may occur on Ribier Grapes. Do not apply within 3 days of harvest.

LADINO CLOVER, VETCH (FOR SEED): Aphids, Spider Mites, Leafhoppers—1½ to 2 pts. in water to cover 1 acre.

Chevron Chemical Company

Ortho Division/San Francisco, Calif. 94119

Richmond, California Fresno, California

Des Moines, Iowa Perth Amboy, New Jersey Orlando, Florida



Form 7732-1 Product 1708 Made in U.S.A.

EPA Reg. No. 239-616-AA



ORTHO

Malathion 5 Emulsive



(INSECTICIDE)

Active Ingredients	By Wt
*Malathion	55%
Aromatic Petroleum Derivative Solvent	32%
Inert Ingredients	13%
*O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate.	
Contains 5 lbs. Malathion per gallon.	



READ THE LABEL

KEEP OUT OF REACH OF CHILDREN

CAUTION: Harmful if swallowed. Avoid breathing spray mist. Avoid contact with skin; wash thoroughly after using. Do not store or transport near feed or food.

Note to Physician: Atropine is antidotal.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

Highly toxic to fish. Do not contaminate any body of water. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

NET CONTENTS 5 GALLONS LIQUID

DIRECTIONS, AND WITH APPLICABLE STATE AND
OR DILUTE INTO FOOD OR DRINK CONTAINERS.

AMOUNT IN 100 GALS. OF WATER UNLESS OTHER-

WATER PER ACRE UNLESS OTHERWISE NOTED:
ton—3 to 20 gals. Deciduous Fruit Crops—60 to
minimum of 15 gals. water per acre).
Orchard Crops—10 to 20 gals.

specified. In Orchards—Codling Moth, Oriental Fruit
after until controlled.

gs, Leafhoppers, Spider Mites, Alfalfa Weevil Larvae,
over. Armyworms—2 pts. per acre in water to cover.
e not working in the field or are not hanging on

ints per acre in water to cover. Do not use during
e in 20 to 30 gals. water by ground application or

loth, Forbes Scale, Woolly Aphid—1 pt. Tent Cater-
Apple Aphid, Rosy Apple Aphid—1½ pts. Malathion
d within 4 weeks of harvest. Do not apply within 3

les—2 pts. per acre in water to cover. Do not apply

2¾ pts. per acre. Aphids, Cucumber Beetles, Leaf-

it.

when first berries turn blue and at 10-day intervals

GREENS, TURNIPS: Aphids, Imported Cabbage-
acre, making full coverage application every 7 days.
s Sprouts, Cauliflower, Cabbage, Kale and Mustard

within 7 days of harvest.

acre. Loopers (on lettuce)—3 pts. per acre. Do not
leaf Lettuce within 14 days of harvest.

occur on certain varieties of sweet cherries, particu-

Scale, Yellow Scale, Purple Scale, Black Scale (sin-
Purple Scale—Use 1 qt. per 100 gals. water. Thrips
oom. Do not apply within 7 days of harvest. Do not

n water to cover. Do not apply to clover in bloom.

otton Leafworm, Cotton Leafworm, Cotton Leaf Per-
z—1 to 2 qts. in water to cover 1 acre. Repeat as

anberry Fruitworm—1½ pints per acre. Do not ap-

rms—1½ pts. per acre. Do not apply to cucurbits
y to stems and vines at base of plants. Do not
umpkins within 3 days of harvest.

t apply within 3 days of harvest.

ulfured Molasses per acre in water to cover. Do not

e-Spotted Bud Moth—1 pt. May be used up to har-


r on Ribier Grapes. Do not apply within 3 days of

½ to 2 pts. in water to cover 1 acre.



ORTHO

Malathion 5 Emulsive



(INSECTICIDE)

Active Ingredients	By Wt.
*Malathion	55%
Aromatic Petroleum Derivative Solvent	32%
Inert Ingredients	13%
*O,O-dimethyl dithiophosphate of diethyl me. aptosuccinate.	
Contains 5 lbs. Malathion per gallon.	



READ THE LABEL

KEEP OUT OF REACH OF CHILDREN

CAUTION: Harmful if swallowed. Avoid breathing spray mist. Avoid contact with skin; wash thoroughly after using. Do not store or transport near feed or food. **Note to Physician:** Atropine is antidotal. **DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.**

Highly toxic to fish. Do not contaminate any body of water. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

NET CONTENTS 5 GALLONS LIQUID

OKRA: Japanese Beetles—2 pts. per acre in water to cover. Do not apply after the pods 4 days. Do not apply to Green onions within 3 days of harvest.

ORNAMENTALS: Aphids, Spider Mites, Whitefly, Mealybugs, Thrips, Japanese Beetle, For Leafhoppers, Scurfy Scale, European Pine Shoot Moth—1½ pts. Soft Brown Scale, Mo Juniper Scale—2½ pts. Oystershell Scale, Lacebugs—1 pt. Pine Needle Scale—4 pts. p after first. Oak Kermes, Fletcher Scale, Florida Red Scale, Bagworms, Azalea Scale, A Miners, Tent Caterpillar—2 pts. Euonymus Scale—1 to 1½ pts. **To Control Listed Scale** at time scale crawlers have settled on foliage. Do not use on Boston, Maidenhair and P **PEACHES:** Oriental Fruit Moth, Plum Curculio, Spider Mites—2 pts. Do not apply with **PECANS:** Spider Mites, Aphids—1½ to 2 pts. May be applied up to harvest.

PEARS: Pear Psylla, Aphids, Woolly Aphids, Mealybugs, Spider Mites—1 to 2 pts. Codlin 2 pts. Injury may occur under certain conditions in the Northwest on Bosc Pears. Do not **PEAS:** Pea Aphid—1½ pts. per acre as needed. Pea Weevil—1 quart in 5 to 10 gals. days of harvest or 7 days if forage is to be fed to livestock.

PEPPERMINT, SPEARMINT: Flea Beetle Adults—1½ pints per acre in water to cover. Do **PEPPERS:** Aphids—1 pt. Pepper Maggots—2½ pts. per acre. Do not apply within 3 da **PLUMS, PRUNES:** Mealy Plum Aphid—1 pt. Do not apply within 3 days of harvest.

POTATOES: Aphids, Leafhoppers—1 pt. per acre. May be used up to harvest. **RASPBERRIES:** Aphids, Rose Scale—1½ qts. per 100 gals. water. Do not apply within **SMALL GRAINS (Barley, Oats, Rye, Wheat in grain forms):** Young Grasshoppers—1½ pt hatching areas when nymphs are young. Do not apply within 7 days of harvest.

STRAWBERRIES: Aphids, Spider Mites, Lygus Bugs, Spittlebugs, Potato Leafhoppers, berry Root Weevil—1½ pts. Field Crickets—1½ to 3 pts. per acre. Strawberry Root We ing and work into top 6 to 8 inches of soil. Do not apply within 3 days of harvest.

CORN (Maize for grain or forage): Young Grasshoppers—1½ pts. per acre in water to c occur in the whorl and silk stage, using this type malathion product. Do not apply withi

TOBACCO: Aphids—1 to 1½ pts. in water to cover 1 acre. Make full-coverage applicati **TOMATOES:** Aphids—1 pt. per acre. One application. Spider Mites—1½ pts. per acre. apply within 1 day of harvest.

WALNUTS: Aphids, Red Spider—1 to 1½ pts. Repeat at 7 to 10 days for Red Spider. M **STORED GRAIN**

For a Residual Wall, Floor and Machinery Spray in Grain Elevators, in Treating Truck Beds ing grain—Use 1 gal. to 25 gals. water making thorough coverage application, contacti ing places. Before applying spray, clean elevators, box cars, etc. thoroughly. Remove a respirator. Wear protective clothing to minimize skin exposure.

For the Protection of Stored Grains: Wheat, Oats, Rice, Corn, Rye, Barley, Grain Sorghu Weevil, Granary Weevil, Saw-toothed Grain Beetle, Flat Grain Beetle, Red Flour Be Borer, Cereal Leaf Beetle—Apply at rate of 1 pt. in 2 to 5 gals. water per 1,000 bushel turned into final storage. Indian Meal Moth—Apply at rate of ½ pt. in 1 to 2 gals. surface area. Apply to spray evenly over the surface of the clean or uninfested grain. A into storage and repeat if necessary.


NOTE: ORTHO Malathion 5 Emulsive may be used under the requirements of the Khap diesel oil emulsions are indicated, as prescribed by the current quarantine instructio **LIVESTOCK**

Do not treat calves under 1 month of age. Do not use on lactating dairy animals. Do within 2 weeks of freshening. Apply complete coverage spray. One treatment may be s contamination of feed and food products, also drinking fountains and feed troughs. Av ment and water. **BEEF CATTLE: Lice**—1 gal. per 100 gals. water. **Ticks**—1 to 2 gals. p **VICES:** To reduce lice, use 7½ Tablespoonfuls or 3¼ fl. ozs. (2½ actual Malathion) i time before use. There may also be a reduction in horn flies. These devices should be ma 35 to 45 head of cattle and re-treated every 2 to 3 weeks. Do not apply or make accessi per 100 gals. water. Apply complete coverage also to pens and litter. **SHEEP, GOATS:** Li water. Repeat application after 2 to 3 weeks if needed. Do not treat animals under 1 m **Northern Fowl Mite, Poultry Lice, and Chicken Red Mite** (as a supplement to premise ti water per 100 to 150 birds. Repeat in 4 to 6 weeks or when necessary. **POULTRY HOUSI Mite, Poultry Lice, Flies**—4 Tablespoonfuls per gal. water. **Poultry Ticks**—6 to 7 ozs. | Apply spray liberally to walls, ceilings and adjacent areas. Force spray into cracks and for surface sprays. **ROOST PAINT: Chicken Red Mite, Poultry Lice**—4 Tablespoonfuls p Brush on at rate of 1 pt. per 150 feet of roost.

CONDITIONS OF SALE: 1. Chevron Chemical Company (Chevron) warrants that this n scription on the label and is reasonably fit for use as directed hereon. Chevron neither representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guaraplied, concerning this material.

2. Critical and unforeseeable factors beyond Chevron's control prevent it from eliminatir of chemicals. Such risks include, but are not limited to, damage to plants and crops, complete control, and damage caused by drift to other plants or crops. Such risks occur fit for the uses stated hereon and even though label directions are followed. Buyer a risks and liability (except those assumed by Chevron under 1 above) resulting from har

THO Malathion 5 Emulsive



ICIDE)

By Wt.
55%
Petroleum Derivative Solvent 32%
..... 13%
Dithiophosphate of diethyl mercaptosuccinate.
Contains 5 lbs. Malathion per gallon.

READ THE LABEL

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid breathing spray mist. Avoid contact thoroughly after using. Do not store or transport near feed or food. Atropine is antidotal. DO NOT DRINK, SPILL OR STORE NEAR HEAT OR OPEN FLAME. Do not apply under conditions that may cause possible drift to food, forage or other plantings that might be damaged. Do not apply to crops thereof rendered unfit for sale, use or consumption. Do not contaminate any body of water. Shrimp and crab may be affected at rates recommended on this label. Do not apply where these are likely to occur. Do not apply where runoff is likely to occur. Do not apply when wind is likely to favor drift from areas treated. Do not contaminate water by cleaning equipment. Dispose of wastes. Apply this product only as specified on this label. Malathion is highly toxic to bees exposed to direct treatment or residues on crops. For more information, contact your Cooperative Agricultural Extension Office.

CONTENTS 5 GALLONS LIQUID

OKRA: Japanese Beetles—2 pts. per acre in water to cover. Do not apply after the pods begin to form.

ONIONS: Onion Thrips—1½ pts. in water to cover 1 acre. **Onion Maggots**—2½ pts. per acre when flies appear. Repeat every 4 days. Do not apply to Green onions within 3 days of harvest.

ORNAMENTALS: Aphids, Spider Mites, Whitefly, Mealybugs, Thrips, Japanese Beetle, Four-Lined Leaf Bug, Tarnished Plant Bug, Leafhoppers, Scurfy Scale, European Pine Shoot Moth—1½ pts. **Soft Brown Scale, Monterey Pine Scale, Black Scale Crawlers, Juniper Scale**—2½ pts. **Oystershell Scale, Lacebugs**—1 pt. **Pine Needle Scale**—4 pts. per acre. Make second application 2 days after first. **Oak Kermes, Fletcher Scale, Florida Red Scale, Bagworms, Azalea Scale, Magnolia Scale, Pine Leaf Scale, Leaf Miners, Tent Caterpillar**—2 pts. **Euonymus Scale**—1 to 1½ pts. **To Control Listed Scale**—Apply a thorough, full-coverage spray at time scale crawlers have settled on foliage. Do not use on Boston, Maidenhair and Pteris ferns, and some species of Crassula.

PEACHES: Oriental Fruit Moth, Plum Curculio, Spider Mites—2 pts. Do not apply within 7 days of harvest.

PECANS: Spider Mites, Aphids—1½ to 2 pts. May be applied up to harvest.

PEARS: Pear Psylla, Aphids, Woolly Aphids, Mealybugs, Spider Mites—1 to 2 pts. **Codling Moth, Plum Curculio, Leaf Rollers**—2 pts. Injury may occur under certain conditions in the Northwest on Bosc Pears. Do not apply within 1 day of harvest.

PEAS: Pea Aphid—1½ pts. per acre as needed. **Pea Weevil**—1 quart in 5 to 10 gals. water per acre. Do not apply within 3 days of harvest or 7 days if forage is to be fed to livestock.

PEPPERMINT, SPEARMINT: Flea Beetle Adults—1½ pints per acre in water to cover. Do not apply within 7 days of harvest.

PEPPERS: Aphids—1 pt. **Pepper Maggots**—2½ pts. per acre. Do not apply within 3 days of harvest.

PLUMS, PRUNES: Mealy Plum Aphid—1 pt. Do not apply within 3 days of harvest.

POTATOES: Aphids, Leafhoppers—1 pt. per acre. May be used up to harvest.

RASPBERRIES: Aphids, Rose Scale—1½ qts. per 100 gals. water. Do not apply within 1 day of harvest.

SMALL GRAINS (Barley, Oats, Rye, Wheat in grain forms): Young Grasshoppers—1½ pints per acre in water to cover. Apply to hatching areas when nymphs are young. Do not apply within 7 days of harvest.

STRAWBERRIES: Aphids, Spider Mites, Lygus Bugs, Spittlebugs, Potato Leafhoppers, Strawberry Leafroller, Whiteflies, Strawberry Root Weevil—1½ pts. **Field Crickets**—1½ to 3 pts. per acre. **Strawberry Root Weevil**—Apply to soil surface before planting and work into top 6 to 8 inches of soil. Do not apply within 3 days of harvest.

CORN (Maize for grain or forage): Young Grasshoppers—1½ pts. per acre in water to cover when nymphs are young. Injury may occur in the whorl and silk stage, using this type malathion product. Do not apply within 5 days of harvest or forage.

TOBACCO: Aphids—1 to 1½ pts. in water to cover 1 acre. Make full-coverage application. Repeat as necessary.

TOMATOES: Aphids—1 pt. per acre. One application. **Spider Mites**—1½ pts. per acre. **Drosophila**—2½ pts. per acre. Do not apply within 1 day of harvest.

WALNUTS: Aphids, Red Spider—1 to 1½ pts. Repeat at 7 to 10 days for Red Spider. May be applied up to harvest.

STORED GRAIN
For a Residual Wall, Floor and Machinery Spray in Grain Elevators, in Treating Truck Beds, Box Cars and Ships' Holds before loading grain—Use 1 gal. to 25 gals. water making thorough coverage application, contacting all cracks, corners, crevices and hiding places. Before applying spray, clean elevators, box cars, etc. thoroughly. Remove and burn all sweepings and debris. Wear respirator. Wear protective clothing to minimize skin exposure.

For the Protection of Stored Grains: Wheat, Oats, Rice, Corn, Rye, Barley, Grain Sorghum against Confused Flour Beetle, Rice Weevil, Granary Weevil, Saw-toothed Grain Beetle, Flat Grain Beetle, Red Flour Beetle, Rusty Grain Beetle, Lesser Grain Borer, Cereal Leaf Beetle—Apply at rate of 1 pt. in 2 to 5 gals. water per 1,000 bushels. Apply as the grain is being loaded or turned into final storage. Indian Meal Moth—Apply at rate of ½ pt. in 1 to 2 gals. of water per 1,000 square feet of grain surface area. Apply to spray evenly over the surface of the clean or uninfested grain. Apply immediately after grain is loaded into storage and repeat if necessary.

NOTE: ORTHO Malathion 5 Emulsive may be used under the requirements of the Khapra Beetle quarantine, where water or diesel oil emulsions are indicated, as prescribed by the current quarantine instructions.

LIVESTOCK
Do not treat calves under 1 month of age. Do not use on lactating dairy animals. Do not use on non-lactating dairy animals within 2 weeks of freshening. Apply complete coverage spray. One treatment may be sufficient. Repeat only if needed. Avoid contamination of feed and food products, also drinking fountains and feed troughs. Avoid contamination of milk, milk equipment and water. **BEEF CATTLE: Lice**—1 gal. per 100 gals. water. **Ticks**—1 to 2 gals. per 100 gals. water. **BACK-RUBBING DEVICES:** To reduce lice, use 7½ Tablespoonfuls or 3¼ fl. ozs. (2% actual Malathion) in 1 gal. fuel oil. Mix thoroughly each time before use. There may also be a reduction in horn flies. These devices should be made continuously accessible, one to each 35 to 45 head of cattle and re-treated every 2 to 3 weeks. Do not apply or make accessible to dairy animals. **HOGS: Lice**—1 gal. per 100 gals. water. Apply complete coverage also to pens and litter. **SHEEP, GOATS: Lice, Ticks, Keds**—1 gal. per 100 gals. water. Repeat application after 2 to 3 weeks if needed. Do not treat animals under 1 month old. **POULTRY (Direct application): Northern Fowl Mite, Poultry Lice, and Chicken Red Mite** (as a supplement to premise treatments)—2 Tablespoonfuls per gal. water per 100 to 150 birds. Repeat in 4 to 6 weeks or when necessary. **POULTRY HOUSES: Northern Fowl Mite, Chicken Red Mite, Poultry Lice, Flies**—4 Tablespoonfuls per gal. water. **Poultry Ticks**—6 to 7 ozs. per 1 gal. water. Repeat as necessary. Apply spray liberally to walls, ceilings and adjacent areas. Force spray into cracks and crevices. Use a high pressure sprayer for surface sprays. **ROOST PAINT: Chicken Red Mite, Poultry Lice**—4 Tablespoonfuls per 1 gal. water. Repeat as necessary. Brush on at rate of 1 pt. per 150 feet of roost.

CONDITIONS OF SALE: 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.
2. Critical and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Chevron under 1 above) resulting from handling, storage and use of this material.