



# Karathane\* 1 Dust

[FUNGICIDE]

NET WEIGHT 50 LBS.

Active Ingredients	By Wt.
2,4-Dinitro-5-octyl** phenyl crotonate	0.73%
2,6-Dinitro-4-octyl** phenyl crotonate	
Nitrooctyl** phenols principally dinitro	0.05%
Inert Ingredients	99.22%

\*\*A mixture of 1-methylheptyl, 1-ethylhexyl, and 1-propylpentyl isomers

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER.

BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE

This product is for use only as directed on the label by ground or air application.

### DIRECTIONS

Cover plants thoroughly directing dust to underside of leaves also. Apply when mildew first appears. This product is suitable for application by power duster, hand duster or airplane

**CUCUMBERS, CANTALOUPE, SQUASH: Powdery Mildew**—30 to 40 lbs. per acre. Apply at first sign of disease. Repeat at 1 to 2 week intervals as necessary to maintain control. Do not apply within 7 days of harvest.



READ THE LABEL

KEEP OUT OF REACH OF CHILDREN

**CAUTION:** May cause skin irritation or sensitization. Avoid skin contact and inhalation of dust. In case of contact, wash thoroughly. Keep out of eyes. Do not take internally. Do not store or transport near feed or food. **Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.**

Highly toxic to fish. Do not contaminate any body of water. Do not contaminate water by cleaning of equipment, or disposal of wastes.

**FLOWERS (Roses, Aster, Larkspur, Chrysanthemums, Delphinium, Calendula, Begonia, Belladonna, Dahlia, Gerbera (Transvaal Daisy), Antirrhinum, Hydrangea, African Violet (Saintpaulia), Snapdragon): Powdery Mildew**—For greenhouse and outdoor plants, apply a light uniform coverage of dust to all parts of plants. Repeat as necessary. Do not use on roses when temperatures exceed 90° F.

**GRAPES: Powdery Mildew**—50 pounds per acre. Apply at first sign of disease. Repeat at 1 to 2-week intervals as necessary to maintain control. Do not apply within 21 days of harvest.


**NURSERY STOCK (Apples): Powdery Mildew**—Apply 25 to 30 lbs. per acre. Repeat at weekly to 10-day intervals as necessary.

**ORNAMENTAL SHRUBS (Hydrangea, Evonymus): Powdery Mildew**—Apply a uniform coverage of dust to all parts of plants. Repeat at weekly to 10-day intervals as necessary.

**CONDITIONS OF SALE 1.** Chevron Chemical Company Chevron warrants that this material conforms to the chemical description on the label and is reasonably fit for use as a fungicide. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or MERCHANTABILITY, guaranteed or implied, concerning this material.

**2.** Chevron does not warrant or assume any liability beyond Chevron's duty to prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks are not covered though the product is reasonably fit for the use intended herein and even though label directions are followed. Buyer and user acknowledge and assume all risks of liability except those assumed by Chevron under 1 above in connection with handling, storage, and use of this material.

© 1977 Chevron Chemical Company, Pat. Off. U.S., Perm. & Mex. Co.



## CHEVRON CHEMICAL COMPANY

FORM 4329 X

EPA Reg. No. 239-848

ORTHO DIVISION - San Francisco, California 94119

Product 2149

Made in U. S. A.

RICHMOND, CALIFORNIA

HONOLULU, HAWAII

FRESNO, CALIFORNIA

DES MOINES, IOWA

PERTH AMBOY, NEW JERSEY

ORLANDO, FLORIDA

