



[INSECTICIDE]

Perthane*Malathion 10-4 Dust

NET WEIGHT 50 LBS.

Active Ingredients	By Wt
Diethyl Diphenyl Dichloroethane	2.1
Related Compounds	1.2
Malathion**	40.0
Inert Ingredients	86.0%
**O,O dimethyl dithiophosphate of diethyl mercaptosuccinate	

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PEST CIDE IN ORIGINAL CONTAINER.

This product is formulated for application by ground or air equipment.

DIRECTIONS

Use 1.5 pounds per acre, unless otherwise specified. Thorough coverage is essential. Repeat application as necessary.

CABBAGE, BRUSSELS SPROUTS, CAULIFLOWER:
Cabbage Worm, Cabbage Looper, Aphids, Thrips
Do not apply within 7 days of harvest.



READ THE LABEL

KEEP OUT OF REACH OF CHILDREN

CAUTION:

Do not apply under conditions involving possible drift to food-bearing plants that might be damaged or the crop thereby rendered unfit for use, unless the crop is protected by a protective netting or other suitable means. Do not apply to plants that are highly toxic to man or animals. Do not apply to plants that are highly toxic to man or animals. Do not apply to plants that are highly toxic to man or animals. Do not apply to plants that are highly toxic to man or animals.

LETTUCE: Cabbage Looper, Leafhoppers, Imported Cabbage Worm, Aphids, Thrips—Apply at weekly to 10-day intervals. Do not apply within 7 days of harvest and to leaf lettuce within 14 days of harvest.

SPINACH: Leafhoppers, Cabbage Looper, Aphids, Thrips—Apply at weekly to 10-day intervals. Do not apply within 7 days of harvest.

CONDITIONS

CHEVRON CHEMICAL COMPANY

Circle 5, 517

USDA Reg. No. 239-1022

ORTHO DIVISION — San Francisco, California 94120

Circle 2617

Made in U.S.A.

RICHMOND, CALIFORNIA

PORTLAND, OREGON

FRESNO, CALIFORNIA



DES MOINES, IOWA

CHERRY HILL, NEW JERSEY

HONOLULU, HAWAII

ORLANDO, FLORIDA