

DIRECTIONS FOR USE

READ INSTRUCTIONS CAREFULLY. DO NOT MIX OR APPLY WITH OTHER CHEMICALS UNLESS SPECIFICALLY RECOMMENDED. For best results, apply PHALITAN Rose & Garden Fungicide in water as indicated below. For best results, apply with a BIRTH SPRAYETTE or pump sprayer. To use in a designated amount of PHALITAN Rose & Garden Fungicide in water, mix the water to make a thick slurry paste then add the required amount of water. Agitate during application. Use water at a minimum of 100°F to allow it to start.

Plant	Disease	Application
Apple	Black Spot	Apply 1/2 to 1/4 teaspoon per gallon of water weekly during periods of rapid new growth and severe infection and twice weekly during periods of rapid new growth and severe infection caused by cool humid rains or high humidity.
Blueberry	Leaf Spot	Apply 1/2 to 1/4 teaspoon per gallon of water weekly during periods of rapid new growth and severe infection and twice weekly during periods of rapid new growth and severe infection caused by cool humid rains or high humidity.
Camellia	Powdery Mildew	Apply 1/2 to 1/4 teaspoon per gallon of water weekly during periods of rapid new growth and severe infection and twice weekly during periods of rapid new growth and severe infection caused by cool humid rains or high humidity.
Chrysanthemum	Powdery Mildew	Apply 1/2 to 1/4 teaspoon per gallon of water weekly during periods of rapid new growth and severe infection and twice weekly during periods of rapid new growth and severe infection caused by cool humid rains or high humidity.
Citrus	Citrus Scab	Apply 1/2 to 1/4 teaspoon per gallon of water weekly during periods of rapid new growth and severe infection and twice weekly during periods of rapid new growth and severe infection caused by cool humid rains or high humidity.
Flowering Quince	Black Spot	Apply 1/2 to 1/4 teaspoon per gallon of water weekly during periods of rapid new growth and severe infection and twice weekly during periods of rapid new growth and severe infection caused by cool humid rains or high humidity.
Hydrangea	Leaf Spot	Apply 1/2 to 1/4 teaspoon per gallon of water weekly during periods of rapid new growth and severe infection and twice weekly during periods of rapid new growth and severe infection caused by cool humid rains or high humidity.
Japanese Quince	Black Spot	Apply 1/2 to 1/4 teaspoon per gallon of water weekly during periods of rapid new growth and severe infection and twice weekly during periods of rapid new growth and severe infection caused by cool humid rains or high humidity.
Japanese Spindle	Black Spot	Apply 1/2 to 1/4 teaspoon per gallon of water weekly during periods of rapid new growth and severe infection and twice weekly during periods of rapid new growth and severe infection caused by cool humid rains or high humidity.
Japanese Yew	Black Spot	Apply 1/2 to 1/4 teaspoon per gallon of water weekly during periods of rapid new growth and severe infection and twice weekly during periods of rapid new growth and severe infection caused by cool humid rains or high humidity.
Juniper	Leaf Spot	Apply 1/2 to 1/4 teaspoon per gallon of water weekly during periods of rapid new growth and severe infection and twice weekly during periods of rapid new growth and severe infection caused by cool humid rains or high humidity.
Lily	Black Spot	Apply 1/2 to 1/4 teaspoon per gallon of water weekly during periods of rapid new growth and severe infection and twice weekly during periods of rapid new growth and severe infection caused by cool humid rains or high humidity.
Magnolia	Black Spot	Apply 1/2 to 1/4 teaspoon per gallon of water weekly during periods of rapid new growth and severe infection and twice weekly during periods of rapid new growth and severe infection caused by cool humid rains or high humidity.
Manzanita	Black Spot	Apply 1/2 to 1/4 teaspoon per gallon of water weekly during periods of rapid new growth and severe infection and twice weekly during periods of rapid new growth and severe infection caused by cool humid rains or high humidity.
Myrtle	Black Spot	Apply 1/2 to 1/4 teaspoon per gallon of water weekly during periods of rapid new growth and severe infection and twice weekly during periods of rapid new growth and severe infection caused by cool humid rains or high humidity.
Nandina	Black Spot	Apply 1/2 to 1/4 teaspoon per gallon of water weekly during periods of rapid new growth and severe infection and twice weekly during periods of rapid new growth and severe infection caused by cool humid rains or high humidity.
Osage Orange	Black Spot	Apply 1/2 to 1/4 teaspoon per gallon of water weekly during periods of rapid new growth and severe infection and twice weekly during periods of rapid new growth and severe infection caused by cool humid rains or high humidity.
Pine	Needle Cast	Apply 1/2 to 1/4 teaspoon per gallon of water weekly during periods of rapid new growth and severe infection and twice weekly during periods of rapid new growth and severe infection caused by cool humid rains or high humidity.
Prunella	Black Spot	Apply 1/2 to 1/4 teaspoon per gallon of water weekly during periods of rapid new growth and severe infection and twice weekly during periods of rapid new growth and severe infection caused by cool humid rains or high humidity.
Rose	Black Spot, Powdery Mildew	Apply 1/2 to 1/4 teaspoon per gallon of water weekly during periods of rapid new growth and severe infection and twice weekly during periods of rapid new growth and severe infection caused by cool humid rains or high humidity.
Rose of Sharon	Black Spot	Apply 1/2 to 1/4 teaspoon per gallon of water weekly during periods of rapid new growth and severe infection and twice weekly during periods of rapid new growth and severe infection caused by cool humid rains or high humidity.
Syringa	Black Spot	Apply 1/2 to 1/4 teaspoon per gallon of water weekly during periods of rapid new growth and severe infection and twice weekly during periods of rapid new growth and severe infection caused by cool humid rains or high humidity.
Taxus	Needle Cast	Apply 1/2 to 1/4 teaspoon per gallon of water weekly during periods of rapid new growth and severe infection and twice weekly during periods of rapid new growth and severe infection caused by cool humid rains or high humidity.
Wisteria	Black Spot	Apply 1/2 to 1/4 teaspoon per gallon of water weekly during periods of rapid new growth and severe infection and twice weekly during periods of rapid new growth and severe infection caused by cool humid rains or high humidity.

CITRUS Citrus Scab - Use 1/2 to 1/4 teaspoon per gallon of water weekly during periods of rapid new growth and severe infection and twice weekly during periods of rapid new growth and severe infection caused by cool humid rains or high humidity.

COMBINATION INSECT AND DISEASE CONTROL SPRAY FOR INSECT CONTROL AND LISTED DISEASES: Add 1 Teaspoon ORTHO Insect Spray or 1 to 1 1/2 Teaspoons of ORTHO Insect Spray depending on what insects are present for each quart of diluted spray. Always mix diluted amount PHALITAN Rose & Garden Fungicide in water to form a thick slurry paste before final dilution and addition of insecticide. Do not use combination on citrus. **FOR ROSES** - Apply fungicide weekly during spring and fall in new seasons. During summer months, apply PHALITAN Rose & Garden Fungicide at regular intervals for disease control. Add ORTHO Insect Spray or ORTHO Insect Spray only when necessary for insect control.

Do not apply more than two consecutive applications of ORTHO Insect Spray in combination with PHALITAN Rose & Garden Fungicide. Do not spray during periods of extremely high temperatures (90°F) as they may occur.

FOLIAGE FEEDING PLUS Insect and Disease Control: To foliage feed plants twice a month add ORTHO Rose & Garden Food B 12-4 at 2 teaspoons per gallon to PHALITAN Rose & Garden Fungicide or the above combination. Note caution on product labels.

COMPATIBILITY: PHALITAN Rose & Garden Fungicide is compatible with most insecticides and miticides. Always refer to product labels for specific information.

CAUTION: Avoid contact with eyes, nose, mouth, and skin. Wash thoroughly with soap and water if contact occurs. Do not eat, drink, or smoke while using this product.

Note to Physicians: This product contains a chemical which is known to be a carcinogen. It may cause cancer in humans. It may also cause birth defects or other reproductive problems. Avoid contact with eyes, nose, mouth, and skin. Wash thoroughly with soap and water if contact occurs. Do not eat, drink, or smoke while using this product.

STORAGE AND DISPOSAL: Store in original container. Do not use if container is damaged. Do not use if contents are solid, lumpy, or otherwise unusable.

NOTICE: This product is not for use on lawns, golf courses, or other areas where children or pets may play. Do not use on lawns, golf courses, or other areas where children or pets may play.



Chevron Chemical Company
 10000 Westheimer Road
 Houston, Texas 77042
 (713) 465-1000



Chevron

ORTHO

PHALTAN Rose & Garden Fungicide



A Spray for Control of
 Black Spot, Powdery Mildew,
 and Certain Other Plant Diseases



COPY A

Add 2-3 Tablespoonfuls of ISOTOX Insect Killer or 2-3 Tablespoonfuls of ORTHENE Systemic Insect Control, depending on what insects are present, for each gallon of diluted spray.

COPY B

ORTHENE Systemic Insect Control

CONFIDENTIAL
PROPERTY OF
CHEVRON CHEMICAL COMPANY
UNLESS ACCEPTED BY EPA