

Evron

ORTHO

TRITHION 4 Flowable

(INSECTICIDE)

Active ingredient	By Wt.
Carbophenothion**	43%
Inert ingredients	57%

**S-[p-chlorophenylthio) methyl] O,O-diethyl phosphorodithioate
Contains — 4 Pounds Trithion Per Gallon
U. S. Pat. No. 2,793,224.



DO NOT TAKE INTERNALLY
DO NOT INHALE
DO NOT GET ON SKIN
KEEP OUT OF REACH OF CHILDREN

WARNING:

MAY BE FATAL IF SWALLOWED. Do not get in eyes, on skin or clothing. In case of contact, wash immediately with soap and water; for eyes, flush with water for 15 minutes. Wash hands, arms and face thoroughly with soap and water before eating or smoking. For commercial application, wear rubber gloves, protective clothing, rubber overshoes and goggles. Wash contaminated clothing with soap and hot water before re-use. Do not store or transport near feed or food. TRITHION may cause cholinesterase inhibition. ATROPINE IS ANTIDOTAL. 2-PAM is also antidotal and may be administered in conjunction with atropine. Do not apply or allow to drift to areas occupied by unprotected humans, beneficial animals or onto adjoining food and forage crops about to be harvested. Do not apply oil sprays to trees where soil moisture is low or where soil moisture has recently been low. Do not apply oil sprays when temperatures are in excess of 90° F. or in periods of low humidity caused by dry winds. Do not combine oil sprays with sulfur sprays or apply within 30 days following a sulfur spray. Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

Toxic to fish and wildlife. Birds feeding on treated areas may be killed. Do not contaminate any body of water nor apply to any area not specified on the label.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

*T.M. Reg. Stauffer Chemical Company for insecticide-acaricide.

NET CONTENTS 5 GALLONS LIQUID

[Faint, mostly illegible text, likely a list of pests or crops treated by the product.]

[Faint, mostly illegible text, likely a list of pests or crops treated by the product.]

ORTHO

TRITHION 4 Flowable

(INSECTICIDE)

Active Ingredient	By Wt.
Carbophenothion**	43%
Inert Ingredients	57%
**S-(p-chlorophenylthio) methyl O,O-diethyl phosphorodithioate	
Contains — 4 Pounds Trithion Per Gallon	
U. S. Pat. No. 2,793,224.	



DO NOT TAKE INTERNALLY
DO NOT INHALE
DO NOT GET ON SKIN
KEEP OUT OF REACH OF CHILDREN

WARNING: MAY BE FATAL IF SWALLOWED. Do not get in eyes, on skin or clothing. In case of contact, wash immediately with soap and water; for eyes, flush with water for 15 minutes. Wash hands, arms and face thoroughly with soap and water before eating or smoking. For commercial application, wear rubber gloves, protective clothing, rubber overshoes and goggles. Wash contaminated clothing with soap and hot water before re-use. Do not store or transport near feed or food. TRITHION may cause cholinesterase inhibition. ATROPINE IS ANTIDOTAL. 2-PAM is also antidotal and may be administered in conjunction with atropine. Do not apply or allow to drift to areas occupied by unprotected humans, beneficial animals or onto adjoining food and forage crops about to be harvested. Do not apply oil sprays to trees where soil moisture is low or where soil moisture has recently been low. Do not apply oil sprays when temperatures are in excess of 90° F. or in periods of low humidity caused by dry winds. Do not combine oil sprays with sulfur sprays or apply within 30 days following a sulfur spray. Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

Toxic to fish and wildlife. Birds feeding on treated areas may be killed. Do not contaminate any body of water nor apply to any area not specified on the label.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

*T.M. Reg. Stauffer Chemical Company for insecticide acaricide

NET CONTENTS 5 GALLONS LIQUID



ACCEPTED

ORTHO TRITHION* 4 Flowable

25872-4

E STATE AND
CONTAINERS

BE NOTED:
60 to

phid, Woolly
ut not more
R DELAYED
ng Mite and

gals. water.
No. Overwin.
X Oil Spray.
rithin 7 days

h—1/2 pint.
1/2 to 1 pint
TRITHION 4

Leathopper,
graze or cut
p residue to

—1 pint per
thin 21 days

pts. per acre

ps—1/2 pint
phid Eggs—
1 of 1 quart


itrus Mite—
ricola Scale,
s ORTHOL-K
y within 30

low 30 days
rithion may

es, Cinnabar
ids. Do not

2 pts per

2 pints per



FIGS: Two Spotted Mites—1/2 to 1 pt. per acre in water to cover. Do not apply within 7 days of harvest.

GARDEN BEETS: Bean Aphid, Serpentine Leaf Miner, Two-Spotted Mites, Potato Flea Beetle—1 to 2 pints per acre in water to cover. Do not apply within 21 days of harvest. Do not graze or feed crop residue to livestock.

GRAPES: Grape Leafhopper, Two-Spotted Mites—1/2 pint per acre in sufficient water for coverage. Do not apply within 30 days of harvest. Do not apply to sensitive varieties such as Charbono, Gamay, and Muscat.

OLIVE: Olive Scale, San Jose Scale, Black Scale, Oleander Scale—1/2 to 1 pint plus 1 to 2 gallons ORTHOL-K Flowable Light Medium or ORTHOL-K READY-MIX Light Medium. Apply a minimum of 1 quart TRITHION 4 Flowable per acre. Do not apply within 60 days of harvest.

ONION: From Seeds, Sets or Transplants (for dry bulb or green)—Onion Maggot Control: Furrow Treatment, Muck Soil: 1 1/2 to 2 qts. per acre. Mineral Soil: 1 qt. to 1 1/2 qts. per acre. Apply at time of seedling or transplanting by spraying in the planting furrow. Mix in 100 gallons of water and apply 3 gals. of mixture per 1000 ft. of furrow. Keep the mixture agitated.

ORNAMENTALS: Camellias, Carnations, Gleditsias, Evergreen and Deciduous Shrubs; Two-Spotted Mites, Spruce Spider Mite, Monterey-Pine Aphid, Rose Aphid, Soft Scale Insects—1/2 to 1 1/2 pints.

PEACHES, NECTARINES: Summer Spray: European Red Mites, Brown Almond Mites, Two-Spotted Mites—1/2 to 1 pint per 100 gallons of water but use no more than 2 lbs. actual Trithion per acre. 1 quart per acre in 20 gallons of water by air application. Do not apply within 30 days of harvest. DORMANT SPRAYS: San Jose Scale, European Fruit Lecanium, Olive Scale—1/2 to 1 pt. plus 1 to 2 gallons of KLEENUP Flowable Emulsion or KLEENUP READY-MIX Oil Spray. Apply a minimum of 1 quart TRITHION 4 Flowable per acre. Do not apply within 30 days of harvest.

PEAS: Pea Aphid, Serpentine Leaf Miner, Two-Spotted Mites—1 to 1 1/2 pints per acre in water to cover. Do not apply within 7 days of harvest. Do not graze or feed crop residue to livestock.

PECAN: Two-Spotted Mites—1/2 pint. Apply a minimum of 1 quart TRITHION 4 Flowable per acre. Do not apply after husks open. Do not graze livestock on cover crops grown in treated pecan groves.

PEPPER, PIMENTO: Serpentine Leaf Miner, Potato Flea Beetle, Potato Leafhopper, Tropical Mite, Two-Spotted Mites—1 to 2 pints per acre in water to cover. Do not apply within 7 days of harvest.

PLUM, PRUNE: SUMMER SPRAYS: Mealy Plum Aphid, Leaf Curl Plum Aphid, Leafhoppers, Two-Spotted Mites, Clover (Brown Almond) Mite, European Red Mite—1/2 pint. DORMANT SPRAYS: Olive Scale, European Fruit Lecanium, San Jose Scale, Overwintering Mite and Aphid Eggs—1/2 to 1 pint plus 1 to 2 gallons KLEENUP READY-MIX Oil Spray or KLEENUP Flowable Emulsion. Apply a minimum of 1 quart TRITHION 4 Flowable per acre. Do not apply within 30 days of harvest.

POST-HARVEST APPLICATION TO DECIDUOUS FRUIT TREES (Listed on Label): To decrease overwintering populations of Aphids, Leafhoppers, Spider Mites—1/2 to 1 pint. Apply a minimum of 1 quart TRITHION 4 Flowable per acre. For control of San Jose Scale, European Fruit Lecanium, Aphid and Mite Eggs—1/2 to 1 pint plus 1 to 2 gallons of ORTHOL-K Flowable Light Medium or ORTHOL-K READY-MIX Oil Spray. Cover crops should not be grazed or fed to livestock.

SEED CROPS (Alfalfa, Clover, Beans, Peas, Sugar Beets): Brown Wheat Mite, Bean Aphid, Pea Aphid, Spotted Alfalfa Aphid, Serpentine Leaf Miner, Two-Spotted Mites—1 to 2 pints per acre in water to cover. Do not feed crop residue to livestock. Cover crops should not be grazed or fed to livestock.

SORGHUM: Sorghum Midge, Two-Spotted Mites—1 pt. per acre in water to cover. Do not make more than 2 applications per season. Do not graze or cut forage within 21 days after application.

SPINACH, BEET TOPS: Serpentine Leaf Miner, Two-Spotted Mites, Bean Aphid—1/2 to 1 pint per acre in water to cover. Do not apply within 21 days of harvest.

STRAWBERRY: Two-Spotted Mites, Strawberry Aphid—1 pint per acre in water to cover. Do not apply within 3 days of harvest.

SUGAR BEET: Bean Aphid, Serpentine Leaf Miner, Potato Flea Beetle, Two-Spotted Mites—1 to 2 pints per acre in water to cover. Do not apply within 14 days of harvest.

TOMATO: Serpentine Leaf Miner, Tomato Russet Mite, Two-Spotted Mites, Tropical Mite, Potato Flea Beetle, Potato Leafhopper, Potato Psyllid—1 to 2 pints per acre in water to cover. Do not apply within 7 days of harvest.

WALNUTS: SUMMER: European Red Mite, Two-Spotted Mites, Walnut Aphid, Codling Moth and for suppression of European Fruit Lecanium—1 to 2 quarts per acre in water to cover. Best results for scale insect suppression are obtained if application made after eggs have hatched. DORMANT: Calico Scale, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 quarts per acre plus 2 gallons KLEENUP Flowable Emulsion or KLEENUP READY-MIX Oil Spray in water to cover. Do not apply after husks have opened. Do not apply more than 2 gals. of oil per acre. Do not graze livestock on cover crops grown in treated walnut groves.

MIXING DIRECTIONS: BEFORE USING AGITATE CONTENTS BY SHAKING OR ROLLING CONTAINER. Pour recommended amount of this material into partially filled spray tank. Add balance of water to fill tank. Keep agitator running during filling, mixing and spraying operations. Do not allow mixture to stand. If mixture does not mix readily, but tends to separate as an oily layer, do not use as injury to plants may result. Do not combine emulsifiable liquids with wettable powders in the same spray tank unless previous use of the materials being mixed has proven them compatible.

CONDITIONS OF SALE: 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though the directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Chevron under 1. above) resulting from handling, storage, and use of this material.