


TRITHION[®] 25 Wettable



ORTHO

ACCEPTED

MAY 10 1973

U.S. DEPARTMENT OF AGRICULTURE
OFFICE OF THE FEDERAL INSPECTOR
FOR PESTICIDE REGULATION
WASHINGTON, D.C. 20250

(INSECTICIDE)

Active Ingredient: Carbophenothion^{††} 25%
Inert Ingredients: 75%
†† S-(1-chloro-2-methyl-2-propyl) O,O-dimethyl phosphorodithioate
U.S. Pat. No. 2,793,224

READ ENTIRE LABEL USE STRICTLY IN ACCORDANCE WITH WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS

COMPLETELY EMPTY BAG AND DISPOSE OF WASTE PESTICIDE BY BURYING, BURN BAG IMMEDIATELY AND STAY OUT OF SMOKE

USE SPECIFIED AMOUNTS IN 100 GALS WATER UNLESS OTHERWISE SPECIFIED

1. Orchard: Codling Moth, Oriental Fruit Moth and Plum Curculio

APPLE, CRAB APPLE, PEAR, QUINCE, APRICOT, PEACH, NECTARINES, PLUMS, PRUNES, CHEERY, SUMMER SPRAYS: Aphids, Apple, Rosy Apple Wasp, Apple Leaf Curl, Plum, Mealy Plum, Spider Mites, Leafhopper, Leghopper, Cherry Rust Mites

As a DORMANT SPRAY on Apples and Pears and as a DORMANT SPRAY on Other Listed Crops: Scale and Overwintering Aphid and Mite Eggs

ALMOND, SUMMER SPRAYS: Aphids, Spider Mites, Leafhoppers
As a DORMANT SPRAY: Scale and Overwintering Aphid and Mite Eggs

POST-HARVEST APPLICATION TO DECIDUOUS FRUIT TREES: To Decrease Overwintering Populations of Aphids, Leafhoppers, Spider Mites—1 to 2 lbs. per acre in 4 to 10 gal. water for control of Scale, Aphid and Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water of a Superior Type Oil. Cover crops completely before harvest.

BEANS (Shelled): Bean Aphid, Mexican Bean Beetle, Serpentine Leaf Miner, Potato Leafhopper, Two Spotted Mite. 2 lbs. per acre in water to cover. For Forest Beans, within 21 days of harvest, spray with 20 lbs. per acre of Trifluralin to control weeds. Do not graze or feed crop residue to livestock.

*T.M. Reg. Stauffer Chemical Company for insecticide acaricide

NET WEIGHT 50 LBS.

Chevron Chemical Company

Ortho Division/San Francisco, Calif. 94119

Richmond, California Fresno, California

Des Moines, Iowa Perth Amboy, New Jersey Orlando, Florida

Form 5487-L Product 2645 Made in U.S.A.

EPA Reg. No. 239-1055-AA

SNAP BEANS, LIMA BEAN, SOYBEAN, in pods: Bean Aphid, Mexican Bean Beetle, Serpentine Leaf Miner, Potato Leafhopper, Two Spotted Mite

CANTALOUPE, WATERMELONS: Melon Aphid, Serpentine Leaf Miner, Potato Leafhopper, Two Spotted Mite

CITRUS: Citrus Red (Purple) Mite, Citrus Rust Mite, Yucca Mite, Two Spotted Mite, Texas Citrus Mite

CUCUMBERS: Citrus Red (Purple) Mite, Citrus Rust Mite, Yucca Mite, Two Spotted Mite, Texas Citrus Mite

CUCUMBERS, SUMMER SQUASH: Melon Aphid, Potato Leafhopper, Serpentine

Leaf Miner, Two Spotted Mite—2 to 4 lbs. per acre in water to cover. Do not apply within 7 days of harvest.

EGGPLANT, PEPPER, PIMENTO: Serpentine Leaf Miner, Two Spotted Mites, Tropical Mite, Potato Flea Beetle, Potato Leafhopper—2 to 4 lbs. per acre in water to cover. Do not apply within 7 days of harvest.

BEETS, GARDEN: Bean Aphid, Potato Flea Beetle, Serpentine Leaf Miner, Two Spotted Mite

GRAPE: Grape Leafhopper, Two Spotted Mite—2 lbs. per acre in water to cover. Do not apply within 7 days of harvest.

OLIVE: Olive Scale, Scale, Scale

ONIONS: Greenhouse Bulb Onion Maggot, Furrow Treatment—4 to 8 lbs. Methocel Seed Coating Treatment—1 lb. per acre in 100 gal. water. Slurry Onion Seed Treatment—1 lb. per 25 gal. water.

STRAWBERRY: Two Spotted Mite, Strawberry Aphid—2 lbs. per acre in water to cover.

SUGAR BEET: Bean Aphid, Serpentine Leaf Miner, Potato Flea Beetle, Two Spotted Mite—2 to 4 lbs. per acre in water to cover. Do not apply within 14 days of harvest.

WALNUT, SUMMER: European Red Mite, Two Spotted Mite, Walnut Aphid, Codling Moth and for suppression of European Fruit Lecanium—4 to 6 lbs. per acre in 4 to 10 gal. water. Walnut Husk Fly Control (California Only)—Use 4 lbs. per acre in 100 gal. water. Do not apply within 7 days of harvest.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

STOP

DO NOT TAKE INTERNALLY
DO NOT INHALE
DO NOT GET ON SKIN

KEEP OUT OF REACH OF CHILDREN

WARNING: MAY BE FATAL IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN. Avoid breathing dust and spray mist. Wear protective clothing. Do not eat, drink, or smoke while using this material. Immediately wash contaminated clothing and water exposed skin and face thoroughly with soap and water. Wash all contaminated clothing with soap and water before reuse. Do not store or transport near food, feed, or other animal products. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, ponds and streams. Do not apply to any area where it is toxic to fish and wildlife.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

BEST DOCUMENT AVAILABLE

Leaf Miner, Two Spotted Mite—2 to 4 lbs. per acre in water to cover. Do not apply within 7 days of harvest.

EGGPLANT, PEPPER, PIMENTO: Serpentine Leaf Miner, Two Spotted Mites, Tropical Mite, Potato Flea Beetle, Potato Leafhopper—2 to 4 lbs. per acre in water to cover. Do not apply within 7 days of harvest.

BEETS, GARDEN: Bean Aphid, Potato Flea Beetle, Serpentine Leaf Miner, Two Spotted Mite

GRAPE: Grape Leafhopper, Two Spotted Mite—2 lbs. per acre in water to cover. Do not apply within 7 days of harvest.

OLIVE: Olive Scale, Scale, Scale

ONIONS: Greenhouse Bulb Onion Maggot, Furrow Treatment—4 to 8 lbs. Methocel Seed Coating Treatment—1 lb. per acre in 100 gal. water. Slurry Onion Seed Treatment—1 lb. per 25 gal. water.

STRAWBERRY: Two Spotted Mite, Strawberry Aphid—2 lbs. per acre in water to cover.

SUGAR BEET: Bean Aphid, Serpentine Leaf Miner, Potato Flea Beetle, Two Spotted Mite—2 to 4 lbs. per acre in water to cover. Do not apply within 14 days of harvest.

WALNUT, SUMMER: European Red Mite, Two Spotted Mite, Walnut Aphid, Codling Moth and for suppression of European Fruit Lecanium—4 to 6 lbs. per acre in 4 to 10 gal. water. Walnut Husk Fly Control (California Only)—Use 4 lbs. per acre in 100 gal. water. Do not apply within 7 days of harvest.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

As a DORMANT SPRAY: European Red Mite, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs—1 to 2 lbs. per acre in 4 to 10 gal. water. Do not apply more than once per year.

TRITHION®*

Wettable

(INSECTICIDE)

Active Ingredient
Carbophenothion**

By Wt.
25%
75%

Inert Ingredients

S 1,1-dichlorophenylthio methyl O,O-diethyl phosphorodithioate.

U. S. Pat. No. 2,793,224

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

EMPTY BAG IMMEDIATELY AND STAY OUT OF SMOKE.

USE SPECIFIED AMOUNTS IN 100 GALS. WATER UNLESS OTHERWISE SPECIFIED.

Apple, Crab Apple, Pear, Quince, Oriental Fruit, etc. (see label for Curculio)

APPLE, CRAB APPLE, PEAR, QUINCE, APRICOT, PEACH, NECTARINES, PLUMS, PRUNES, CHERRY: SUMMER SPRAYS Aphids, Apple, Rosy Apple, Woolly, Apple, Leaf Curl Plum, Mealy Plum, Spider Mites, Codling Moth, Leafrollers, Cherry Rust Mites

... DORMANT ... DORMANT ...

COMPLETELY EMPTY BAG AND DISPOSE OF WASTE PROPERLY BY BURNING.
BURN BAG IMMEDIATELY AND STAY OUT OF SMOKE.

Do not store in or around the home.

USE SPECIFIED AMOUNTS IN 100 GALS. WATER UNLESS OTHERWISE SPECIFIED. Make thorough coverage spray. Application should be made only by a licensed person using approved safety equipment. In Orchards—Codling Moth, Oriental Fruit Moth and Plum Curculio Apply at petal fall and every 10 to 14 days thereafter until controlled.

APPLE, CRAB APPLE, PEAR, QUINCE, APRICOT, PEACH, NECTARINES, PLUMS, PRUNES, CHERRY: SUMMER SPRAYS—Aphids, Apple, Rosy Apple, Woolly Apple, Leaf Curl Plum, Mealy Plum, Spider Mites, Codling Moth, Leafhoppers, Cherry Rust Mites—1 lb. plus 3 to 4 lbs. of a Superior Type Oil per acre. Apply to Apples, Crab Apples, Pears and Quinces only after petal fall and every 10 to 14 days until harvest. As a DORMANT or DELAYED DORMANT SPRAY on Apples and Pears and as a DORMANT SPRAY on Other Listed Crops: Scale and Overwintering Aphid and Mite Eggs—1 to 2 lbs. ORTHO TRITHION 25 Wettable plus 3 to 1 1/2 gals. of a Superior Type Oil with thorough coverage.

ALMOND: SUMMER SPRAYS—Aphids, Spider Mites, Leafhoppers 1 lb. Use at least 4 lbs. but not more than 10 lbs. per acre. If almond hull are to be used for food, do not apply within 7 days of harvest. As a DORMANT SPRAY: Scale and Overwintering Aphid and Mite Eggs—1 to 2 lbs. ORTHO TRITHION 25 Wettable plus 3 to 1 1/2 gals. of a Superior Type Oil with thorough coverage.

POST-HARVEST APPLICATION TO DECIDUOUS FRUIT TREES: To Decrease Overwintering Populations of Aphids, Leafhoppers, Spider Mites—1 to 2 lbs. (Minimum—4 lbs. per acre.) For control of Scale, Aphid and Mite Eggs—1 to 2 lbs. plus 3 to 1 1/2 gals. of a Superior type Oil. Cover crops should not be grazed or fed to live stock.

BEANS (Shelled): Bean Aphid, Mexican Bean Beetle, Serpentine Leaf Miner, Potato Leafhopper, Two-Spotted Mites—2 lbs. per acre in water to cover. Do not apply more than twice per season. Do not graze, cut for forage or harvest beans within 21 days following the last application. Do not graze or feed crop residue to live stock.

*T.M. Reg. Stauffer Chemical Company for insecticide-acaricide

NET WEIGHT 50 LBS.

Chevron Chemical Company

Ortho Division/San Francisco, Calif. 94119

Richmond, California

Fresno, California

Des Moines, Iowa

Perth Amboy, New Jersey

Orlando, Florida

Form 5487-L

Product 2645

Made in U.S.A.

EPA Reg. No. 239-1055-AA

ORTHOK Flowable Light Medium. Allow 30 days between applications. Do not apply within 30 days of harvest. Do not apply to kumquat or citrus.

CITRUS (Florida): Citrus Red (Purple) Mite, Texas Citrus Mite, Six-Spotted Mite—1 to 2 lbs. Apply when needed. Allow 30 days between applications. Do not apply within 14 days of harvest, 30 days when combined with oil spray. Caution: TRITHION may cause injury to grapefruit when applied to stem or fall.

COTTON: Aphid, Leaf Perforator, Fleahopper, Leafworm, Tropical Mite, Two-Spotted Mites, Cinnabar Mite—2 to 4 lbs. per acre in water to cover. Minimum—4 lbs. per acre for control of cotton leaf perforator and aphids. Do not graze or feed animals in treated fields.

CUCUMBERS, SUMMER SQUASH: Melon Aphid, Potato Leafhopper, Serpentine



DO NOT TAKE INTERNALLY

DO NOT INHALE

DO NOT GET ON SKIN

KEEP OUT OF REACH OF CHILDREN

WARNING:

MAY BE FATAL IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN. Avoid breathing dust and spray mist. Wear dust separator and goggles. Do not get in eyes, on skin or on clothing. In case of contact, immediately remove contaminated clothing and wash skin thoroughly with soap and water; for eyes, flush with water for 15 minutes. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use. Do not store or transport near feed or food. TRITHION may cause cholinesterase inhibition. Atropine is Antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine. Do not apply or allow to drift to areas occupied by unprotected human, beneficial animals or onto adjoining food and forage crops about to be harvested. Do not apply oil sprays to trees when soil moisture is low or when soil moisture has recently been low. Do not apply oil spray when temperatures are in excess of 90° F. or in periods of low humidity caused by dry wind. Do not combine oil sprays with sulfur sprays or apply within 30 days following a sulfur spray. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, ponds and streams. Do not apply to any area not specified on the label.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

25



ORTHO

ACCEPTED

MAY 10 1973

UNDER THE FEDERAL INSECTICIDE ACT AND FEDERAL FOOD, DRUG AND COSMETIC ACT REGISTERED UNDER NO. 100-100000 SUBJECT TO ATTACHED COMMENTS.

SNAP BEANS, LIMA BEAN, SOYBEAN (in pods): Bean Aphid, Mexican Bean Beetle, Serpentine Leaf Miner, Potato Leafhopper, Two-Spotted Mites—2 to 3 lbs. per acre in water to cover. Do not apply more than twice per season. Do not graze or cut vines for hay within 21 days after application. Do not apply within 7 days of harvest. Do not graze or feed crop residue to livestock.

CANTALOUPE, WATERMELONS: Melon Aphid, Serpentine Leaf Miner, Potato Flea Beetle, Two-Spotted Mites—2 to 4 lbs. per acre in water to cover. Do not apply within 5 days of harvest.

CITRUS (California, South): Citrus Red (Purple) Mite, Citrus Rust Mite, Yuma Mite, Six-Spotted Mite, Texas Citrus Mite—1 to 2 lbs. Allow 30 days between applications. Do not apply within 14 days of harvest. California Red Scale, Citricola Scale, Cottony Cushion Scale, Florida Red Scale, Purple Scale, Soft Scale, Yellow Scale—2 to 3 lbs. ORTHO TRITHION 25% Emulsion, 1.5 to 2.0 lbs. ORTHO LK READY-MIX Light, Medium or Heavy Duty Emulsion. Allow 30 days between applications.

CITRUS (Florida): Citrus Red (Purple) Mite, Texas Citrus Mite, Six-Spotted Mite—1 to 2 lbs. Apply when needed. Allow 30 days between applications. Do not apply within 14 days of harvest, 30 days when combined with other citrus insecticides. TRITHION emulsion may be applied to citrus trees and shrubs.

COTTON: Aphid, Leaf Perforator, Fleahopper, Leafworm, Tropical Mite, Two-Spotted Mites, Cinnabar Mite—2 to 4 lbs. per acre in water to cover. Mite—4 lbs. per acre for control of cotton leaf perforator and aphids. Do not re-apply to newly emerged insects in treated fields.

CUCUMBERS, SUMMER SQUASH: Melon Aphid, Potato Leafhopper, Serpentine

Leaf Miner, Two-Spotted Mites—2 to 4 lbs. per acre in water to cover. Do not apply within 7 days of harvest.

EGGPLANT, PEPPER, PIMENTO: Serpentine Leaf Miner, Two-Spotted Mites, Tropical Mite, Potato Flea Beetle, Potato Leafhopper—2 to 4 lbs. per acre in water to cover. Do not apply within 7 days of harvest.

BEETS, GARDEN: Bean Aphid, Potato Flea Beetle, Serpentine Leaf Miner, Two-Spotted Mites—1 to 2 lbs. Do not apply within 21 days of harvest. Do not graze or feed to livestock.

GRAPES: Grape Leafhopper, Two-Spotted Mites—2 lbs. per acre in water to cover. Do not apply within 30 days of harvest. Do not apply to sensitive varieties such as Concord, Conroy, and Niagara.

OLIVE: Olive Scale, San Jose Scale—1 lb. per acre in water to cover. Do not apply within 30 days of harvest. Do not apply to sensitive varieties such as Mission, Black Leaf, and California Black Leaf.

ONIONS (Green or Dry Bulb): Onion Maggots—Furrow Treatment—4 to 8 lbs. per acre in water to cover. Do not apply within 30 days of harvest. Methocel Seed Coating Treatment—1 lb. per 100 lbs. seed. Do not apply to onions that are heavily coated. U.S. 4 lb. ORTHO TRITHION 25% Emulsion per 25 lbs. seed. Do not apply to onions that are heavily coated. Slurry Onions—Seed Treatment—1 lb. per 100 lbs. seed. Do not apply to onions that are heavily coated.

STRAWBERRY: Two Spotted Mites, Strawberry Aphid—2 lbs. per acre in water to cover. Do not apply within 30 days of harvest.

SUGAR BEET: Bean Aphid, Serpentine Leaf Miner, Potato Flea Beetle, Two-Spotted Mite—2 to 4 lbs. per acre in water to cover. Do not apply within 30 days of harvest.



DO NOT TAKE INTERNALLY

DO NOT INHALE

Active ingredient: Ortho-Trithion 25% Emulsion. Contains 25% active ingredient. Do not use in residential areas. For more information, contact your local agricultural agent.

CITRUS (Florida): Citrus Red (Purple) Mite, Texas Citrus Mite, Six-Spotted Mite—1 to 2 lbs. Apply when needed. Allow 30 days between applications. Do not apply within 14 days of harvest, 30 days when combined with oil spray. Caution: TRITHION may cause injury to grapefruit when applied in summer or fall.

COTTON: Aphid, Leaf Perforator, Fleahopper, Leafworm, Tropical Mite, Two-Spotted Mites, Cinnabar Mite—2 to 4 lbs. per acre in water to cover. (For mites—4 lbs. per acre for control of cotton leaf perforator and aphids.) Do not graze dairy or meat animals in treated fields.

CUCUMBERS, SUMMER SQUASH: Melon Aphid, Potato Leafhopper, Serpentine

plus 1 to 2 gals. ORTHOL-K Flowable Light Medium or ORTHOL-K READY-MIX Light Medium. (Minimum—4 lbs. per acre.) Do not apply within 60 days of harvest.

ONIONS (Green or Dry Bulb): Onion Maggots—Furrow Treatment—4 to 8 lbs. per acre depending on soil types. Spray at time of seeding in planting furrow. **Methocel Seed Coating Treatment**—Coat onion seeds with adequate amount of water or barrel until seed is thoroughly coated. Use 4 lbs. ORTHO TRITHION 25 Wettable per 25 lbs. seed for upland and 1 lb. per 15 lbs. seed for table onions. **Slurry Onion Seed Treatment**—Use 4 lbs. per 25 lbs. onion seed by slurry method.

STRAWBERRY: Two-Spotted Mites, Strawberry Aphid—2 lbs. per acre in water to cover. Do not apply within 3 days of harvest.

SUGAR BEET: Bean Aphid, Serpentine Leaf Miner, Potato Flea Beetle, Two-Spotted Mites—2 to 4 lbs. per acre in water to cover. Do not apply within 14 days of harvest.

WALNUT—SUMMER: European Red Mites, Two-Spotted Mites, Walnut Aphid, Codling Moth and for suppression of European Fruit Lecanium—4 to 8 lbs. per acre in water to cover. Best results for scale insect suppression are obtained if application is made after eggs have hatched. Do not apply after husks have opened. **Walnut Husk Fly Control (California Only)**—Use 4 lbs. plus 2 gal. ORTHO STALEY'S Protein Bait No. 7 in 100 gals. water. Do not apply more than 8 lbs. ORTHO TRITHION 25 Wettable per acre. Apply at first sign of population increase after fly emergence begins in July. Do not apply after husks open. **DORMANT: Calico Scale, European Fruit Lecanium, Frosted Scale, Walnut Aphid and Overwintering Mite Eggs**—4 to 8 lbs. plus 2 gals. KLEENUP Flowable Emulsion or KLEENUP READY-MIX Oil Spray per acre in water for coverage. Do not apply more than 2 gals. of oil per acre. Do not graze livestock on cover crops grown in treated walnut groves.

MIXING DIRECTIONS: Pour recommended amount of this material on surface of water in partially filled spray tank, or pre-mix in a bucket half filled with water and pour into nearly filled spray tank. Add balance of water to fill tank. Keep agitator running during filling, mixing and spraying operations. Do not allow mixture to stand. Do not use in low volume gear-type spray equipment. Do not combine wettable powders with emulsifiable liquids in the same spray tank unless previous use of the materials being combined has proven them compatible.

CONDITIONS OF SALE: 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Chevron under 1 above) resulting from handling, storage, and use of this material.



DO NOT TAKE INTERNALLY

DO NOT INHALE

DO NOT GET ON SKIN

KEEP OUT OF REACH OF CHILDREN

WARNING:

MAY BE FATAL IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN. Avoid breathing dust and spray mist. Wear dust respirator and goggles. Do not get in eyes, on skin or on clothing. In case of contact, immediately remove contaminated clothing and wash skin thoroughly with soap and water; for eyes, flush with water for 15 minutes. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use. Do not store or transport near feed or food. TRITHION may cause cholinesterase inhibition. Atropine is Antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine. Do not apply or allow to drift to areas occupied by unprotected humans, beneficial animals or onto adjoining food and forage crops about to be harvested. Do not apply oil sprays to trees where soil moisture is low or where soil moisture has recently been low. Do not apply oil sprays when temperatures are in excess of 90° F. or in periods of low humidity caused by dry winds. Do not combine oil sprays with sulfur sprays or apply within 30 days following a sulfur spray. **Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.**

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, ponds and streams. Do not apply to any area not specified on the label.

This product is toxic to bees and should not be applied when bees are actively visiting the area.