

# Copper Monohydrate 10 Dust



# ORTHO

**ACCEPTED**  
FEB 11 1974  
UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTER-  
ED UNDER NO. 239-1060-  
TO ATTACHED COMMENTS

## (FUNGICIDE)

Active Ingredient	By Wt.
Copper Sulfate Monohydrate .....	10%
Inert Ingredients .....	90%
Copper expressed as Elemental—3.58%	

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER.

DO NOT REUSE EMPTY CONTAINER. DESTROY IT BY BURYING WITH WASTE OR BURNING. STAY AWAY FROM SMOKE OR FUMES.

### DIRECTIONS

Use 30 to 50 lbs. per acre unless otherwise specified. Best results are obtained by keeping the plants covered during periods of infection. Dust the plants thoroughly. This product is formulated for application by ground or air equipment.

## NET WEIGHT 50 LBS.

### Chevron Chemical Company

Ortho Division/San Francisco, Calif. 94119  
 Richmond, California    Fresno, California  
 Des Moines, Iowa    Perth Amboy, New Jersey    Orlando, Florida  
 Form 4426-P    Product 2187    Made in U.S.A.  
 EPA Reg. No. 239-1060-AA

**CARROTS: Cercospora Blight, Macrosporium Blight**—Begin applications at first sign of disease. Repeat at 7 to 10-day intervals as necessary.

**CELERY: Early Blight, Late Blight and Bacterial Blight**—Apply at weekly intervals in the seedbed and at 7 to 10-day intervals in the field, beginning when plants are established.

**PEARS: Fire Blight (Pear Blight)**—Begin dusting at 15 to 20% bloom, and repeat at 5 to 7-day intervals or before rainy or wet weather.

**PEPPERS, PIMENTOS: Cercospora Leaf Spot, Anthracnose**—Begin applications at first sign of disease. Repeat at 7 to 10-day intervals as necessary.

**POTATOES: Late Blight, Early Blight**—The first application should be made when the plants are 4 to 6 inches high. Repeat at 10-day intervals as necessary.



## READ THE LABEL

KEEP OUT OF REACH OF CHILDREN

**CAUTION:** Harmful if swallowed. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

This product is toxic to fish. Keep out of any body of water. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

**STRAWBERRIES: Leaf Spot (Mycosphaerella)**—Begin applications at first sign of disease. Repeat at 7 to 10-day intervals as necessary.

**TOMATOES: Late Blight, Septoria Leaf Spot and Early Blight**—If disease appears in seedbeds, plants should be dusted at weekly intervals. It is important to dust plants 5 days before or 5 days after transplanting.

**CONDITIONS OF SALE:** 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Chevron under 1 above) resulting from handling, storage, and use of this material.