



ACCEPTED
 Apr 1, 1971
 239-1070
 CHEVRON CHEMICAL COMPANY REGISTERED
 TRADEMARK

Parathion-ZINC COPOSIL® 2-25 Dust

NET WEIGHT 50 LBS.

Active Ingredients	By Wt.
Parathion*	2.0%
Copper (In basic cupric zinc sulfate complex) expressed as metallic	4.5%
Inert Ingredients	93.5%
Zinc (In basic cupric zinc sulfate complex) expressed as metallic	4.5%
*O,O-diethyl O-p-nitrophenyl thiophosphate.	

Use within 30 days of manufacture only, as this product decomposes.

Date of Manufacture _____

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER.

BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE.

DIRECTIONS

Use 20 to 25 pounds per acre unless otherwise specified, depending on the size of plants and insect infestation. Dust listed crops thoroughly before diseases or insects appear. Repeat applications according to local practice. Where disease control is of primary importance closer to harvest, a suitable fungicide should be used alone. This product is designed for application through ground dusters or by airplane. This product is not to be used around homes or home gardens.

BEANS: Bacterial Blight, White Mold, Psyllid, Aphids, Thrips, Red Spider Mites—25 lbs. per acre. Do not apply within 15 days of harvest.

CABBAGE, CAULIFLOWER, BROCCOLI, BRUSSELS SPROUTS, LETTUCE: Aphids, Leaf Miner, Downy Mildew—20 to 25 lbs. per acre. Do not apply to Cabbage within 10 days of harvest and to Cauliflower, Broccoli, Brussels Sprouts and Head Lettuce within 7 days of harvest. Do not apply to Leaf Lettuce within 21 days of harvest.

CELERY: Aphids, Worms, Thrips, Red Spider Mites, Early and Late Blight—Use 20 lbs. per acre. Do not apply within 21 days of harvest. By Airplane—50 lbs. per acre. Do not apply within 30 days of harvest.



POISON



ANTIDOTE: If Swallowed. Give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Have victim lie down and keep quiet. **CALL A PHYSICIAN IMMEDIATELY!** Note to Physicians: Atropine and 2-PAM are Antidotal. If On Skin. In case of contact, remove contaminated clothing and immediately wash skin with soap and water.

DANGER:

KEEP OUT OF REACH OF CHILDREN
 DO NOT TAKE INTERNALLY
 DO NOT INHALE
 DO NOT GET ON SKIN

DANGER:

Poisonous If Swallowed, Inhaled or Absorbed Through Skin! Wear a mask or respirator of a type passed by the U. S. Department of Agriculture for parathion protection. Rapidly Absorbed Through Skin! Do not get in eyes, on skin or on clothing. Wear natural rubber gloves, protective clothing and goggles. In case of contact wash immediately with soap and water. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use. Keep all unprotected persons out of operating areas or vicinity where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated. Do not store or transport near feed or food. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

This product is toxic to fish and wildlife. Keep out of lakes, ponds and streams. Do not apply to any area not specified on the label. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from you. Cooperative Agriculture Extension Service.

CUCUMBERS: Aphids, Pickle and Melon Worms, Red Spider Mites, Blight, Downy and Powdery Mildew—Use 20 lbs. per acre. Apply when diseases appear. Repeat at 7 to 10-day intervals. Do not apply to Cucumbers before vining or within 15 days of harvest.

POTATOES and TOMATOES: Aphids, Leafhoppers, Thrips, Early and Late Blight—Use 22 lbs. per acre. Do not apply to tomatoes within 10 days of harvest and to potatoes within 5 days of harvest.

SQUASH: Aphids, Pickle and Melon Worms, Red Spider Mites, Blight, Downy and Powdery Mildew—17 lbs. per acre. Do not apply within 15 days of harvest.

STRAWBERRIES: Red Spider, Aphids, White Fly, Cyclamen Mite, Thrips, Lygus Bugs, Potato Leafhoppers, Downy and Powdery Mildew—20 lbs. per acre. Repeat against Red Spider in 7 to 10 days. Apply when insects or diseases appear. Do not apply within 14 days of harvest.

WATERMELONS, CANTALOUPE: Aphids, Pickle and Melon Worms, Red Spider Mites, Blight, Downy and Powdery Mildew—Do not apply within 7 days of harvest.

CONDITIONS OF SALE: 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material. 2. Critical and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability except those assumed by Chevron under 1 above) resulting from handling, storage, and use of this material.

CHEVRON CHEMICAL COMPANY

FORM 2997-G

USDA Reg. No. 239-1070

ORTHO DIVISION — San Francisco, California 94120

Product No. 1574

Made in U. S. A.

RICHMOND, CALIFORNIA

PORTLAND, OREGON

FRES CALIFORNIA

DES MOINES, IA

CHERRY HILL, NEW JERSEY

HONOLULU, HAWAII

ORLANDO, FLORIDA

