


Malathion-Sulfur 4-25 Dust

[239-1100]



ORTHO



(INSECTICIDE-FUNGICIDE)

Active Ingredients	By Wt.
Malathion*	4%
Sulfur	25%
Inert Ingredients	71%

*O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER.

BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE.

DIRECTIONS

Make one or more applications as needed, with no applications to food crops later than 3 days before harvest unless otherwise specified. Apply at first sign of insects or disease and repeat at regular intervals as necessary to maintain control. This product is formulated for application by ground or air equipment.

NET WEIGHT 50 LBS.

Chevron Chemical Company

Ortho Division/San Francisco, Calif. 94119

Richmond, California Fresno, California Honolulu, Hawaii
Des Moines, Iowa Perth Amboy, New Jersey Orlando, Florida

Form 3803-S Product 1914 Made in U.S.A.

EPA Reg. No. 239-1100-1AA

APPLES AND PEARS: Aphids, Spider Mites, Woolly Aphid, Pear Psylla, Powdery Mildew—30 to 40 pounds per acre. Do not use sulfur on Apple foliage within 10 days to 2-weeks of an oil spray. This product may injure foliage on some varieties of apples. Sulfur may burn foliage when temperature is excessively high. Do not apply to Pears within 1 day of harvest.

COTTON: Aphids, Two-Spotted Mite, Pacific Mite, Atlantic Mite—25 to 30 pounds per acre.

CUCUMBERS (coastal areas only): Aphids, Spider Mites,

Pickleworm, Cucumber Beetle, Leaf Hoppers, Powdery Mildew—Use 40 lbs. per acre. Do not apply within 1 day of harvest.

GRAPES: Grape Leafhopper, Spider Mites, Powdery Mildew—20 to 30 lbs. per acre. ~~Weedy-bugs~~—25 to 30 lbs. per acre. Apply at first signs of insects or diseases. Repeat at two week intervals or as necessary. May cause injury to foliage on some varieties.

ORNAMENTALS AND NURSERY STOCK: Two-Spotted Mite, Soft Brown Scale, Aphids, Powdery Mildew—30 to 40 pounds per acre. Do not use on Boston, Maidenhair and Pteris ferns, and some species of Crassula.

POTATOES: Aphids, Leafhopper—30 to 40 pounds per acre. May be used up to harvest.

TOMATOES: Russet Mite, Two-Spotted Spider Mite—25 to 35 lbs. per acre. Do not apply within 1 day of harvest. Because of the sulfur content of this dust, first obtain permission from your canner or processor before using it on tomatoes intended for sale to such canner or processor.

CONDITIONS OF SALE: 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability except those assumed by Chevron under 1 above resulting from handling, storage, and use of this material.



READ THE LABEL

KEEP OUT OF REACH OF CHILDREN

CAUTION: Avoid breathing dust. Avoid contact with skin; wash thoroughly after using. Do not store or transport near feed or food. Some fruits and other plants are susceptible to injury from sulfur under certain climatic conditions. The user is advised not to use sulfur on any crop unless local use has proved that sulfur does not damage crops in that locality. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

Highly toxic to fish. Do not contaminate any body of water nor apply when conditions favor drift from the crop area treated.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.