

**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS & DOMESTIC ANIMALS**  
**CAUTION**

**STATEMENT OF PRACTICAL TREATMENT**

Note to Physicians:

**ENVIRONMENTAL HAZARDS**

**DIRECTIONS FOR USE**

**READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.**

**REENTRY STATEMENT**

**STORAGE AND DISPOSAL**

**PROHIBITIONS**

**STORAGE**

**PESTICIDE DISPOSAL**

**CONTAINER DISPOSAL**

**DIRECTIONS**

USE DESIGNATED AMOUNT IN 100 GALS. WATER, UNLESS OTHERWISE SPECIFIED.

**APPLES, CRABAPPLES:** Bitter Rot, Botryospheria, Alternaria Spot, Brooks Fruit Spot, Sooty Blotch, Ploek Rot, Fly Spok, Scab. Apply 2.5 lbs. per acre. Caution: East of the Rocky Mountains, apply at 7 to 14 day intervals.

**AVOCADOS:** Scab. Apply 2.5 lbs. per acre. Begin application when the buds are opening. Apply at 7 to 14 day intervals. Moderate to severe scab may require 2 to 3 weekly treatments as needed.

**BLACKBERRIES, BOYSENBERRIES, DEWBERRIES, LOGANBERRIES, RASPBERRIES:** Fruit Rot, Gray Mold, Spur Blight. 2 lbs. Apply 250 gals. per acre. Apply when berries have set and repeat at 7 day intervals until harvest.

**Chevron**

# ORTHO

## PHALTAN® 50

### Wettable

### Folpet FUNGICIDE

Active Ingredient: Folpet  
 Inert Ingredients: N-[Trichloromethylthio]phthalimide

By Wt.  
 50%  
 50%

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

**NET WEIGHT 40 POUNDS**  
**(18.1 KGS.)**

**Chevron Chemical Company**  
 Ortho Agricultural Chemicals Division  
 San Francisco CA 94120-7144 Richmond CA 94804-0036  
 Product 2632 Made in U.S.A.  
 Form 9716 C EPA Est 239-NJ-1  
 EPA Reg No 239-1102-AA

**BLACKBERRIES, BOYSENBERRIES, DEWBERRIES, LOGANBERRIES, RASPBERRIES:** Fruit Rot, Gray Mold, Spur Blight. 2 lbs. Apply 250 gals. per acre. Apply when berries have set and repeat at 7 day intervals until harvest.

**CHEERY:** Early Blight, Late Blight, Cercospora Leaf Spot. 2 lbs. Apply 250 gals. per acre. Repeat at 7 to 10 day intervals.

**CHERRIES, RED TART:** Leaf Spot, Brown Rot. 2 lbs. Apply 250 gals. per acre. Repeat at 7 to 10 day intervals.

**CITRUS:** Orange, Grapefruit, Lemon, Lime, Tangerine, Melanose, Scale, Green Spot. 2 lbs. Apply 250 gals. per acre. Repeat at 7 to 10 day intervals.

**CRANBERRIES:** Botry, Rot. 2 lbs. Apply 250 gals. per acre. Repeat at 7 to 10 day intervals.

**CUCUMBERS:** Fruit Rot, Rhizoctonia & Pythium. 2 lbs. Apply 250 gals. per acre. Repeat at 7 to 10 day intervals.

**ANTHRACNOSE:** Downy Mildew, Powdery Mildew. 2 lbs. Apply 250 gals. per acre. Repeat at 7 to 10 day intervals.

**GRAPES:** Downy Mildew, Powdery Mildew, Black Rot, Dead Arm. 2 lbs. Apply 250 gals. per acre. Repeat at 7 to 10 day intervals.

**GOOSEBERRIES, CURRANTS:** Leaf Spot. 2 lbs. Apply 250 gals. per acre. Repeat at 7 to 10 day intervals.

**LETTUCE:** Downy Mildew. 2 lbs. Apply 250 gals. per acre. Repeat at 7 to 10 day intervals.

**MELONS, CANTALOUPES, WATERMELON, HONEYDEW, PUMPKINS, SUMMER AND WINTER SQUASH:** Downy Mildew, Powdery Mildew, Anthracnose. 2 lbs. Apply 250 gals. per acre. Repeat at 7 to 10 day intervals.

**ONIONS, GARLIC, LEEKS, SHALLOTS:** Purple Blotch, Downy Mildew. 2 lbs. Apply 250 gals. per acre. Repeat at 7 to 10 day intervals.

**ORNAMENTALS:** Powdery Mildew and Black Spot on ROSES, Septoria Leafspot, Didymella Leafspot on CHRYSANTHEMUMS AND IRIS, Rust and Alternaria Leafspot on CARNATIONS, Alternaria Leafspot, Powdery Mildew and Anthracnose on ZINNIA, MARIGOLDS, CHRYSANTHEMUMS, ASTERS, PHLOX AND SNAPDRAGONS. 2 lbs. Apply 250 gals. per acre. Repeat at 7 to 10 day intervals.

**PYTHIUM AND ROOT ROT ON PANSY, GREENHOUSE:** Damp off on AZALEA Cuttings. 2 lbs. Apply 250 gals. per acre. Repeat at 7 to 10 day intervals.

**TOMATOES:** Anthracnose. 2 lbs. Apply 250 gals. per acre. Repeat at 7 to 10 day intervals.

Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale or consumption.

**COMPATIBILITY:** Folpet is compatible with most other fungicides and insecticides. Do not mix with lime sulfur or other alkaline materials.

**CONDITIONS OF SALE:** This product is sold as a fungicide. It is not a pesticide. It is not to be used for any purpose other than that for which it is sold. It is not to be used in any manner that would be injurious to humans, animals, or the environment. It is not to be used in any manner that would be injurious to crops or other plantings. It is not to be used in any manner that would be injurious to the environment. It is not to be used in any manner that would be injurious to the environment.