



# TEDION<sup>®</sup> 25 Wettable

## [MITICIDE]

### NET WEIGHT 50 LBS.

Active Ingredient	By Wt.
Tetradifon	25%
Inert Ingredients	75%
1,4-dichlorophenyl 2,4,5-trichlorophenyl sulfone	
U. S. Pat. No. 2,812,281	

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

**APPLE, CRABAPPLE, PEAR, QUINCE:** WEST: European Red Mite, Two-Spotted Mite, McDaniel Mite, Canadensis Mite, Carpini Mite, Schoenii Mite. EAST: European Red Mite, Two-Spotted Spider Mite, McDaniel Mite, Canadensis Mite, Carpini Mite, Schoenii Mite—When Oil or Phosphate—Oil Spray. EASTERN AREAS: European Red Mite, Two-Spotted Spider Mite, McDaniel Mite, Canadensis Mite, Carpini Mite, Schoenii Mite—When Oil or Phosphate—Oil Spray.

**APRICOTS, CHERRIES, PEACHES, NECTARINES:** European Red Mite, Two-Spotted Mite, McDaniel Mite — WEST COAST AREAS: 1 lb. per 100 gals. water to cover. EASTERN AREAS: 1 lb. per 100 gals. water to cover.

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION:** Avoid contact with children, domestic animals, feed animals, and birds. Do not use on lawns or ornamentals treated with insecticides. Do not use on plants treated with insecticides. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

**BUSHBERRIES:** Blackberries, Boysenberries, Blueberries, Gooseberries, Loganberries, Raspberries: Two-Spotted Mite—Post-Harvest Treatment Only.

**CITRUS—Grapefruit, Limes, Oranges, Tangelos, Tangerines, Lemons:** Citrus Red Mite, Six-Spotted Mite, Texas Citrus Mite.

**FLORIDA AREAS:** Citrus Red Mite, Six-Spotted Mite, Texas Citrus Mite.

**OTHER AREAS:** Citrus Red Mite, Six-Spotted Mite, Texas Citrus Mite.

**DECIDUOUS FRUITS:** European Red Mite, Two-Spotted Mite, McDaniel Mite, Canadensis Mite, Carpini Mite, Schoenii Mite—When Oil or Phosphate—Oil Spray.

**EVERGREENS:** Spruce Mite—When Oil or Phosphate—Oil Spray.

**FIGS:** Pacific Mite, Two-Spotted Spider Mite—1 lb. to 10 lbs. per acre in water to cover when mites first appear. Do not make more than 2 applications during the fruiting period. Do not apply within 14 days of harvest.

**GRAPES:** Pacific Mite, Two-Spotted Mite, Willamette Mite—4 lbs. per acre in water to cover. Apply when mites first appear and repeat as necessary. Do not make more than 3 applications during the fruiting period. May be applied up to the day of harvest.

**HOPS:** Two-Spotted Mites—1 lb. per 100 gals. water or 4 lbs. per acre in water to cover. Apply when mites first appear. Do not apply more than 2 applications during fruiting period. Do not apply within 14 days of harvest. Do not feed crop refuse to livestock.

**MELONS, PUMPKINS, WINTER SQUASH:** Two-Spotted Mite—2 lbs. per acre in water to cover. Apply when mites appear and repeat as necessary. May be applied up to day of harvest.

**ORNAMENTALS:** Citrus Red Mite, Two-Spotted Mite—1 pound per 100 gals. of water. Spray thoroughly when mites first appear and repeat as required.

**PLUMS, PRUNES:** European Red Mite, Two-Spotted Mite, McDaniel Mite—WEST COAST AREAS: 1 lb. per 100 gals. water or 5 lbs. per acre in water to cover. EASTERN AREAS: 1 lb. per 100 gals. water to cover.

**STRAWBERRIES:** Two-Spotted Mite—4 lbs. per acre in water to cover when mites first appear. Do not make more than 2 applications during the fruiting period. Do not apply within 14 days of harvest.

**TOMATOES:** Tropical Mite, Two-Spotted Mite—1 lb. per 100 gals. of water. Spray thoroughly when mites first appear and repeat as required.