

TRITHION[®] 3 Dust



(INSECTICIDE)

Active Ingredient	By Wt.
Carbophenothion*	3%
Inert Ingredients	97%
*S-[(p-chlorophenylthio) methyl] O,O-diethyl phosphorodithioate.	

U. S. Pat. No. 2,793,224.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER.

BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE.

Apply when air is calm. Repeat as necessary. This product is formulated for application by ground or air equipment. Make thorough coverage of plants with a uniform deposit of dust.

APPLE, CRABAPPLE, PEAR, QUINCE: Apple Aphid, European Red Mite, Rosy Apple Aphid, Two-Spotted Mites, Woolly Apple Aphid, Codling Moth—30 to 40 pounds per acre. Apply at pre-pink stage. Do not apply within 30 days of harvest.

ALMOND, APRICOT, CHERRY, NECTARINE, PEACH, PECAN, PLUM, PRUNE, WALNUT: Aphids, Orchard Mites—40 to 50 pounds per acre. If almond hulls are to be used for feed, do not apply within 7 days of harvest. Do not apply to apricot, cherry, nectarine, peach, plum and prune within 30 days of harvest. Do not use on pecan and walnut after husks open. Do not graze livestock on covercrops grown in treated walnut and pecan groves.

SNAP BEAN, LIMA BEAN, SOYBEAN (in pod): Bean Aphid, Mex-

CITRUS (Grapefruit, Lemons, Limes, Oranges, Tangelos, Tangerines): Citrus Red (Purple) Mite, Citrus Rust Mite, Six-Spotted Mite, Texas Citrus Mite, Yuma Mite—35 to 60 lbs. per acre. Allow 30 days between applications. Do not apply within 14 days of harvest. Do not apply to kumquats or citron.

COTTON: Cotton Aphid, Cotton Leaf Perforator, Cotton Flea Hopper, Cotton Leafworm, Cinnabar Mite, Tropical Mite, Two-Spotted Mites—20 to 30 lbs. per acre. Apply 30 lbs. for control of cotton leaf perforator and aphids. Do not graze dairy or meat animals in treated fields.

CUCUMBER, SUMMER SQUASH: Melon Aphid, Potato Leafhopper, Serpentine Leaf Miner, Two-Spotted Mites—20 to 35 lbs. per acre. Do not apply within 7 days of harvest.

EGGPLANT: Serpentine Leaf Miner, Potato Leafhopper, Potato Flea Beetle, Tropical Mite, Two-Spotted Mite—20 to 35 pounds per acre. Do not apply within 7 days of harvest.

FIGS: Two-Spotted Mites—30 to 40 lbs. per acre. Do not apply within 7 days of harvest.

GARDEN BEETS: Bean Aphid, Leafhoppers, Serpentine Leaf Miner, Two-Spotted Mites, Potato Flea Beetle—15 to 25 pounds per acre.

Do not apply for food crops.

GRAPES: 25 lbs. per acre.

ONIONS: 25 lbs. per acre. Do not apply after planting for suitable growth.

ORNAMENTALS: 25 lbs. per acre. Do not apply to greens, D.

PEA: 25 lbs. per acre. Do not graze or feed to livestock.

PEPPER, P: 25 lbs. per acre. Do not apply to Potato Leaf Miner.

POTATO: 25 lbs. per acre. Do not apply to SEED CROPS.

WHEAT: 25 lbs. per acre. Do not apply to Wheat Mite, Serpentine Mite.

SPINACH: 25 lbs. per acre. Do not apply to Bean Aphid.

STRAWBERRY: 25 lbs. per acre. Do not apply to days of harvest.

SUGAR BEET: 25 lbs. per acre. Do not apply to days of harvest.



READ THE LABEL

KEEP PESTICIDE IN ORIGINAL CONTAINER.

BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE.

Apply when air is calm. Repeat as necessary. This product is formulated for application by ground or air equipment. Make thorough coverage of plants with a uniform deposit of dust.

APPLE, CRABAPPLE, PEAR, QUINCE: Apple Aphid, European Red Mite, Rosy Apple Aphid, Two-Spotted Mites, Woolly Apple Aphid, Codling Moth—30 to 40 pounds per acre. Apply at pre-pink stage. Do not apply within 30 days of harvest.

ALMOND, APRICOT, CHERRY, NECTARINE, PEACH, PECAN, PLUM, PRUNE, WALNUT: Aphids, Orchard Mites—40 to 50 pounds per acre. If almond hulls are to be used for feed, do not apply within 7 days of harvest. Do not apply to apricot, cherry, nectarine, peach, plum and prune within 30 days of harvest. Do not use on pecan and walnut after husks open. Do not graze livestock on cover crops grown in treated walnut and pecan groves.

SNAP BEAN, LIMA BEAN, SOYBEAN (in pod): Bean Aphid, Mexican Bean Beetle, Serpentine Leaf Miner, Potato Leafhopper, Two-Spotted Mites—20 to 25 pounds per acre. Do not apply within 7 days of harvest. Do not graze or feed crop residue to livestock.

BEANS (shelled): Bean Aphid, Mexican Bean Beetle, Serpentine Leaf Miner, Two-Spotted Mites, Potato Leafhopper—20 pounds per acre. Do not apply more than twice per season. Do not graze, cut for forage or harvest beans within 21 days following the last application. Use only on beans to be marketed as shelled dry beans.

CANTALOUPE, WATERMELON: Melon Aphid, Serpentine Leaf Miner, Potato Flea Beetle, Two-Spotted Mites—20 to 35 pounds per acre. Do not apply within 5 days of harvest.

NET WEIGHT 50 LBS.

Chevron Chemical Company

Ortho Division/San Francisco, California 94119

Richmond, California Fresno, California

Des Moines, Iowa Perth Amboy, New Jersey Orlando, Florida

Form 5244-E2 Product 2584 Made in U.S.A.

EPA Reg. No. 239-1187-AA

Serpentine Leaf Miner, Two-Spotted Mites—20 to 35 lbs. per acre. Do not apply within 7 days of harvest.

EGGPLANT: Serpentine Leaf Miner, Potato Leafhopper, Potato Flea Beetle, Tropical Mite, Two-Spotted Mite—20 to 35 pounds per acre. Do not apply within 7 days of harvest.

FIGS: Two-Spotted Mites—30 to 40 lbs. per acre. Do not apply within 7 days of harvest.

GARDEN BEETS: Bean Aphid, Leafhoppers, Serpentine Leaf Miner, Two-Spotted Mites, Potato Flea Beetle—15 to 25 pounds per acre.



READ THE LABEL

KEEP OUT OF REACH OF CHILDREN

CAUTION: HAZARDOUS IF SWALLOWED. Avoid breathing dust. Wear a respirator and goggles. Avoid contact with skin and eyes. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use. Do not store or transport near feed or food. TRITHION may cause cholinesterase inhibition. Atropine is Antidotal. 2-PAM is also antidotal and may be administered in conjunction with Atropine. Do not apply or allow to drift to areas occupied by unprotected humans, beneficial animals or onto adjoining food and forage crops about to be harvested. **Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.**

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, ponds and streams. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Do not apply to any area not specified on the label.

This product is toxic to bees and should not be applied when bees are commonly visiting the area.

PEA: Pe
25 lbs.
graze o
PEPPER,
Potato
lbs. per
SEED C
Wheat
Serpent
acre. D
not be
SPINAC
Bean A
days of
STRAW
pounds
SUGAR
Beetle,
apply w
TOMATO
Beetle, T
Tropical
7 days
CONDIT
warrant
on the
Chevron
tive to
ABILITY,
cerning
2. Critic
prevent
of chem
to plant
complete
or crops
fit for tl
are follo
and liab
resulting
TRITHION
secticide-c

®

3



ORTHO

ACCEPTED

FEB 8 1974

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC INDIAN REGISTRATION
ED UNDER NO. 230-1185

CITRUS (Grapefruit, Lemons, Limes, Oranges, Tangelos, Tangerines): Citrus Red (Purple) Mite, Citrus Rust Mite, Six-Spotted Mite, Texas Citrus Mite, Yuma Mite—35 to 60 lbs. per acre. Allow 30 days between applications. Do not apply within 14 days of harvest. Do not apply to kumquats or citron.

COTTON: Cotton Aphid, Cotton Leaf Perforator, Cotton Flea Hopper, Cotton Leafworm, Cinnabar Mite, Tropical Mite, Two-Spotted Mites—20 to 30 lbs. per acre. Apply 30 lbs. for control of cotton leaf perforator and aphids. Do not graze dairy or meat animals in treated fields.

CUCUMBER, SUMMER SQUASH: Melon Aphid, Potato Leafhopper, Serpentine Leaf Miner, Two-Spotted Mites—20 to 35 lbs. per acre. Do not apply within 7 days of harvest.

EGGPLANT: Serpentine Leaf Miner, Potato Leafhopper, Potato Flea Beetle, Tropical Mite, Two-Spotted Mite—20 to 35 pounds per acre. Do not apply within 7 days of harvest.

FIGS: Two-Spotted Mites—30 to 40 lbs. per acre. Do not apply within 7 days of harvest.

GARDEN BEETS: Bean Aphid, Leafhoppers, Serpentine Leaf Miner, Two-Spotted Mites, Potato Flea Beetle—15 to 25 pounds per acre.

Do not apply within 21 days of harvest if tops are to be used for food or feed.

GRAPES: Grape Leafhopper, Two-Spotted Mites—20 to 25 pounds per acre. Do not apply within 30 days of harvest.

ONIONS (Green or Dry Bulb): Onion Maggot—33 to 66 lbs. per acre depending on soil types. Apply at the time of seeding in the planting furrow, using a combination drill, fertilizer spreader, or suitable granular application equipment.

ORNAMENTALS (Camellias, Carnations, Gladiolus, Violets, Evergreens, Deciduous Shrubs): Spider Mites, Aphids—15 to 25 lbs. per acre.

PEA: Pea Aphid, Serpentine Leaf Miner, Two-Spotted Mites—20 to 25 lbs. per acre. Do not apply within 7 days of harvest. Do not graze or feed crop residue to livestock.

PEPPER, PIMENTO: Serpentine Leaf Miner, Potato Flea Beetle, Potato Leafhopper, Tropical Mite, Two-Spotted Mites—20 to 30 lbs. per acre. Do not apply within 7 days of harvest.

SEED CROPS: (Alfalfa, Clover, Beans, Peas, Sugar Beets): Brown Wheat Mite, Pea Aphid, Bean Aphid, Spotted Alfalfa Aphid, Serpentine Leaf Miner, Two-Spotted Mites—20 to 30 pounds per acre. Do not feed crop residue to livestock. Cover crops should not be grazed or fed to livestock.

SPINACH, BEET TOPS: Serpentine Leaf Miner, Two-Spotted Mites, Bean Aphid—15 to 25 pounds per acre. Do not apply within 21 days of harvest.

STRAWBERRY: Two-Spotted Mites, Strawberry Aphid — 20 to 30 pounds per acre. Do not apply within 3 days of harvest.

SUGAR BEET: Bean Aphid, Serpentine Leaf Miner, Potato Flea Beetle, Two-Spotted Mites—20 to 30 pounds per acre. Do not apply within 14 days of harvest.

By Wt.
..... 3%
..... 97%

CE WITH
LE STATE

SMOKE.
product is
nt. Make
of dust.
ean Red
le Aphid,
ink stage.

PECAN,
0 pounds
not apply
y, nectar-
t. Do not
raze live-
groves.
hid, Mex-
per, Two-



READ THE LABEL

CUCUMBER, SUMMER SQUASH: Melon Aphid, Potato Leafhopper, Serpentine Leaf Miner, Two-Spotted Mites—20 to 35 lbs. per acre. Do not apply within 7 days of harvest.

EGGPLANT: Serpentine Leaf Miner, Potato Leafhopper, Potato Flea Beetle, Tropical Mite, Two-Spotted Mite—20 to 35 pounds per acre. Do not apply within 7 days of harvest.

FIGS: Two-Spotted Mites—30 to 40 lbs. per acre. Do not apply within 7 days of harvest.

GARDEN BEETS: Bean Aphid, Leafhoppers, Serpentine Leaf Miner, Two-Spotted Mites, Potato Flea Beetle—15 to 25 pounds per acre.

per acre.

PEA: Pea Aphid, Serpentine Leaf Miner, Two-Spotted Mites—20 to 25 lbs. per acre. Do not apply within 7 days of harvest. Do not graze or feed crop residue to livestock.

PEPPER, PIMENTO: Serpentine Leaf Miner, Potato Flea Beetle, Potato Leafhopper, Tropical Mite, Two-Spotted Mites—20 to 30 lbs. per acre. Do not apply within 7 days of harvest.

SEED CROPS: (Alfalfa, Clover, Beans, Peas, Sugar Beets): Brown Wheat Mite, Pea Aphid, Bean Aphid, Spotted Alfalfa Aphid, Serpentine Leaf Miner, Two-Spotted Mites—20 to 30 pounds per acre. Do not feed crop residue to livestock. Cover crops should not be grazed or fed to livestock.

SPINACH, BEET TOPS: Serpentine Leaf Miner, Two-Spotted Mites, Bean Aphid—15 to 25 pounds per acre. Do not apply within 21 days of harvest.

STRAWBERRY: Two-Spotted Mites, Strawberry Aphid — 20 to 30 pounds per acre. Do not apply within 3 days of harvest.

SUGAR BEET: Bean Aphid, Serpentine Leaf Miner, Potato Flea Beetle, Two-Spotted Mites—20 to 30 pounds per acre. Do not apply within 14 days of harvest.

TOMATOES: Serpentine Leaf Miner, Potato Psyllid, Potato Flea Beetle, Potato Leafhopper, Tomato Russet Mite, Two-Spotted Mites, Tropical Mite—20 to 30 pounds per acre. Do not apply within 7 days of harvest.

CONDITIONS OF SALE: 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Chevron under 1 above) resulting from handling, storage, and use of this material.

TRITHION® registered trademark of Stauffer Chemical Company for insecticide-acaricide.



READ THE LABEL

KEEP OUT OF REACH OF CHILDREN

CAUTION: HAZARDOUS IF SWALLOWED. Avoid breathing dust. Wear a respirator and goggles. Avoid contact with skin and eyes. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use. Do not store or transport near feed or food. TRITHION may cause cholinesterase inhibition. Atropine is Antidotal. 2-PAM is also antidotal and may be administered in conjunction with Atropine. Do not apply or allow to drift to areas occupied by unprotected humans, beneficial animals or onto adjoining food and forage crops about to be harvested. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, ponds and streams. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Do not apply to any area not specified on the label.

This product is toxic to bees and should not be applied when bees are commonly visiting the area.