

ZINC COPOSIL[®] Sulfur 25-30 Dust



(FUNGICIDE)

Active Ingredients	By Wt.
Copper (in basic cupric zinc sulfate complex) expressed as elemental	4.5%
Sulfur	30.0%
Inert Ingredients	65.5%
Contains 4.5% zinc as elemental derived from basic cupric zinc sulfate complex.	

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP THIS DUST IN ORIGINAL CONTAINER.

EMPTY BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE.

DIRECTIONS

This product is formulated for application by ground or air equipment.

LETTUCE: Downy and Powdery Mildew—40 to 60 lbs. per acre. Apply at first sign of disease and repeat as necessary.

NET WEIGHT 50 LBS.


Chevron Chemical Company

Ortho Division, San Francisco, California 94119

Richmond, California Fresno, California Honolulu, Hawaii
Des Moines, Iowa Perth Amboy, New Jersey Orlando, Florida

Form 2907-B Product 1536 Made in U.S.A.

© 1972 Chevron Chemical Company



239-1235

POTATOES: Late Blight—40 to 60 lbs. per acre. Dust thoroughly and repeat applications at 7-day intervals as necessary. The first application should be made before the disease is expected to appear.


TABLE GRAPE BUNCHES: Bunch Rot (Botrytis)—Dust just after thinning, 25 pounds per acre and a second dusting previous to ripening.

TOMATOES: Late Blight—40 to 60 lbs. per acre. Dust thoroughly before disease appears, usually

starting before early fall rains are expected. Repeat at 7 to 10-day intervals. Because of the sulfur content of this dust, first obtain permission from your canner or processor before using it on tomatoes intended for sale to such canner or processor.

CONDITIONS OF SALE: 1. Chevron Chemical Company Chevron warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability except those assumed by Chevron under 1. above, resulting from handling, storage, and use of this material.



READ THE LABEL

KEEP OUT OF REACH OF CHILDREN

CAUTION: Harmful if swallowed. Some fruits and other plants are susceptible to injury from sulfur under certain climatic conditions. The user is advised not to use sulfur on any crop unless local use has proved that sulfur does not damage crops in that locality. **Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.**

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from areas treated.