

NET CONTENTS

LIQUID

Active Ingredients	By Wt.
Isooctyl ester of 2,4-dichlorophenoxyacetic acid	34.0%
Isooctyl ester of 2,4,5-trichlorophenoxyacetic acid	32.5%
Inert Ingredients	33.5%
Acid Equivalent:	
2,4-dichlorophenoxyacetic acid	22.59%—2 lbs. per gal.
2,4,5-trichlorophenoxyacetic acid	22.59%—2 lbs. per gal.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONTENTS OF CONTAINER IN FOOD OR DRINK CONTAINERS.

DO NOT BURN. NEVER RE-USE

CONTROLS

Aspens
Blackberries, Raspberries and related brambles
Chokecherry
Cottonwood
Coyote Brush
Currants
Elderberry

Elm
Groundcherry
Honeysuckle
Horsenettle
Locust
Maples
Milkvetch
Oaks

Plums
Poison Ivy
Poison Oak
Sassafras
Sumac
Sweetgum
Wild Gooseberries
Wild Grapes
Wild Roses
Willows

And Many Other Weeds

READ THE LABEL

KEEP OUT OF REACH OF CHILDREN

CAUTION

Avoid contact with skin, eyes and clothing. Wash off promptly with plenty of soap and water any material that accumulates on the skin. DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME. Vapors from this product may injure susceptible plants in the immediate vicinity. Avoid spray drift to susceptible plants, as this product may injure such plants as cotton, beans, peas, tomatoes and ornamentals. Coarse sprays are less likely to drift. Apply by airplane, ground rigs and hand dispensers only when there is no hazard from drift. The airplane equipment should have an engine shut-off. High wind velocities should be avoided. Because of the volatility of thoroughly cleaning sprayers and other equipment used with this formulation, it is recommended that such equipment not be used for handling and applying other volatile chemicals. If same equipment is to be used for other purposes, wash thoroughly 2 or 3 times with hot water, kerosene, cleaning soda and strong laundry soap. Do not transport with or store near fertilizers, insecticides, fungicides or seeds. Excessive amounts of this material in the soil may temporarily inhibit seed germination or plant growth. Do not store near foodstuffs. Toxic to fish. Do not contaminate any body of water, by direct application, cleaning of equipment or disposal of wastes and containers.

ACCEPTED

June 11, 1965

DIRECTIONS

Repeat applications may be necessary if regrowth appears. Keep spray tank agitated during application.

FOLIAGE TREATMENT (Along Fences, Roads, Powerlines, Brush Clearing Areas, Pastured Field Areas, Non-Pastured Field Areas, etc.)—3 to 4 quarts per 100 gals. water. Spray to wet thoroughly, when plants are in full leaf, preferably in early summer while making active growth. Spot-treat survivors and seedlings during successive years to control all susceptible species. When present, grasses quickly replace brush and weeds.

BASAL BARK SPRAYING: 4 to 8 gals. per 100 gals. oil (1 to 2 pts. in 3 gals.). Spray the basal to point of run-off 12 to 20 inches around each stem or trunk, also spray the exposed roots and crown area. Apply anytime during the year, however, for selective killing or when susceptible crops are grown nearby, dormant season spraying is preferable. Do not cut trees and shrubs for a full season following treatment, since they die rather slowly. Repeat application may be necessary if new growth appears.

STUMP TREATMENT: 4 to 8 gals. per 100 gals. of oil (1 to 2 pts. in 3 gals.). Spray tops, sides and exposed roots of fresh cut stumps until thoroughly wet, to prevent sprouting.

CONDITIONS OF SALE: 1. Chevron Chemical Company, (Chevron) warrants that this material conforms to the chemical description on the label and is recommended for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guaranteed or representation, express or implied, concerning this material.

2. Chevron disclaims any liability for factors beyond Chevron's control, including all risks in connection with the use of this material. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Chevron under 1 above) resulting from handling, storage, and use of this material.

CHEVRON CHEMICAL COMPANY

FORM 6482 G

USDA Reg. No. 259-1256

Manufactured for

ORTHO DIVISION — San Francisco, California 94120

Product No. 1000

Made in U. S. A.

RICHMOND, CALIFORNIA

PORTLAND, OREGON

FRESNO, CALIFORNIA

DES MOINES, IOWA

CHERRY HILL, NEW JERSEY

DALLAS, TEXAS

HONOLULU, HAWAII

ORLANDO, FLORIDA

