

[239-1288]



Custom Blend No. 5

[INSECTICIDE-FUNGICIDE]

NET WEIGHT 20 LBS.

CONTAINS 25% ORTHOCIDE[®]
AND 47.5% LEAD ARSENATE

Active Ingredients	By Wt.
Captan*	25.0%
Standard Lead Arsenate (PbHAsO ₄)	47.5%
Inert Ingredients	27.5%
*N-(1-trichloromethylthio)-4-cyclohexene-1,2-dicarboximide.	
Lead expressed as Metallic — 28.35%	
Arsenic expressed as Metallic — 10.25%	
Water soluble arsenic, expressed as metallic, not more than—1%	
U. S. Pat. Nos. 2,553,770—2,553,771	

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP IN ORIGINAL CONTAINER UNTIL USED. DO NOT RE-FILL. DO NOT REUSE CONTAINER.

STAKE IN COOL DRY PLACE. PROTECT FROM EXCESSIVE HEAT. BURN EARLY WHEN EMPTY STAKE OUT OF SMOKE.

DIRECTIONS

For use on apples, pears, cherries, and other fruit trees. Apply to the leaves and fruit. Use 4 lbs. per 100 gallons of water. The active ingredient is 25% ORTHOCIDE 50 Wettable and 47.5% LEAD ARSENATE. This product is highly effective against insects and diseases. It is also a fungicide. For use on other fruit trees, see the label directions.

APPLES: Apple Scab, Brook's Fruit Spot, Bitter Rot, Black Rot, Botryosphaeria Rot, Fly-specking, Black Pox, Apple Maggot—4 lbs. per 100 gallons water in petal fall and 3 to 4 cover sprays at 12 to 14 day intervals. Follow the advice of your local agricultural authorities on timing for control of Cherry Fruit Flies. Do not apply to cherries intended for fresh market within 30 days of harvest, or to those intended for processing within 14 days of harvest. Remove residues remaining at harvest by washing, wiping or other effective means.

with comments



DO NOT INHALE
DO NOT GET ON SKIN
DO NOT TAKE INTERNALLY

KEEP OUT OF REACH OF CHILDREN

WARNING: May be Fatal if Swallowed! Avoid contact with skin, eyes, or clothing. Wash thoroughly after using. Avoid breathing dust or spray mist. Avoid contamination of forage and pasture. Avoid contamination of feed or feedstuffs. Keep livestock off treated areas! Foliage injury may occur. Do not apply to Red Delicious, Winesap and other sensitive varieties in early season sprays. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

Disease Control and Apple Maggot—4 lbs. per 100 gallons water. Make applications approximately the first and fifteenth of July or follow your local agricultural authority for timing to control Apple Maggot. In remaining cover sprays use a combination of 1 lb. ORTHOCIDE 50 Wettable plus 1 lb. ORTHO Zineb Wettable for control of summer diseases and include appropriate insecticide. Do not use ORTHO Custom Blend No. 5 within 30 days of harvest and remove excess residue at harvest. The ORTHOCIDE 50 Wettable component may be used up to 7 days after harvest.

SOUR CHERRIES: Cherry Leaf Spot, Plum Curculio, Cherry Fruit Flies, Rose Chafer—4 lbs. per 100 gallons water in petal fall and 3 to 4 cover sprays at 12 to 14 day intervals. Follow the advice of your local agricultural authorities on timing for control of Cherry Fruit Flies. **CAUTION:** Lead Arsenate may injure Man-of-war. Do not apply to cherries intended for fresh market within 30 days of harvest, or to those intended for processing within 14 days of harvest. Remove residues remaining at harvest by washing, wiping or other effective means.

SWEET CHERRIES: Cherry Leaf Spot, Brown Rot, Plum Curculio, Cherry Fruit Flies, Rose Chafer—4 lbs. per 100 gallons in petal fall and 3 to 4 cover sprays at 12 to 14 day intervals and follow the advice of your local agricultural authorities on timing for control of Cherry Fruit Flies. Do not apply to cherries intended for fresh market within 30 days of harvest, or to those intended for processing within 14 days of harvest. Remove residues remaining at harvest by washing, wiping or other effective means.

NOTE: For pre-bloom, bloom and pre-harvest control of Brown Rot; Use 2 lbs. of ORTHOCIDE 50 Wettable. For post-harvest control of Cherry Leaf Spot; Apply 2 lbs. ORTHO PHALTAN 50 Wettable Fungicide per 100 gal. as soon as harvest is complete.

COMPATIBILITY: This product may be combined with most commonly used insecticides, adjuvants and fungicides with the exception of strongly alkaline material such as hydrated lime, Sodium Hydroxide (Caustic), Lime Sulfur and Bordeaux Mixture. It should not be used in combination with or closely following a spray.

CONDITIONS OF SALE: This product is sold as a fungicide, insecticide and acaricide. It is not to be used as a herbicide. It is not to be used on ornamental plants. It is not to be used on lawns. It is not to be used on golf courses. It is not to be used on tennis courts. It is not to be used on swimming pools. It is not to be used on playgrounds. It is not to be used on school grounds. It is not to be used on residential areas. It is not to be used on public places. It is not to be used on private property without the permission of the owner. It is not to be used on any other property without the permission of the owner.

CHEVRON CHEMICAL COMPANY

FORM 5682-E

USDA Reg. No. 239-1288

ORTHO DIVISION — San Francisco, California

Product No. 2809

Made in U.S.A.

RICHMOND, CALIFORNIA

PORTLAND, OREGON

FRESNO, CALIFORNIA

DES MOINES, IOWA

CHERRY HILL, NEW JERSEY

DALLAS, TEXAS

BITTIER, CALIFORNIA

HONOLULU, HAWAII

ATLANTA, GEORGIA

FORT MADISON, IOWA

ORLANDO, FLORIDA