

Zineb 5 Dust

ACCEPTED
AUG 13 1973
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 239-1296 SUBJECT
TO ATTACHED COMMENTS.



ORTHO

(FUNGICIDE)

239-1296

Active Ingredient	By Wt.
Zineb*	5%
Inert Ingredients	95%
Zinc expressed as metallic	1.18%
*Zinc ethylene bisdithiocarbamate	

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER.
BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE.

DIRECTIONS
This product is formulated for application by ground or air equipment.

BEANS: Rust, Anthracnose, Downy Mildew—Use 30 to 40 lbs. per acre. Start dusting immediately upon the appearance of the disease, or when the disease threatens and continue regular applications at 7 to 10-day intervals. **Anthracnose:** Start dusting when the first blossoms appear and continue regular applications at 7 to 10-day intervals. Do not apply within 7 days of harvest.

BROCCOLI, CABBAGE: Downy Mildew—40 to 50 lbs. per acre. Begin application before disease appears. Repeat at 7 to 10-day intervals. Do not apply within 7 days of harvest.

CARROTS: Late Blight—35 to 50 pounds per acre. Apply at first sign of disease and repeat as necessary. Do not apply within 7 days of harvest if treated tops are to be used as food or feed.

NET WEIGHT 50 LBS.

Chevron Chemical Company
Ortho Division, San Francisco, Calif. 94119
Richmond, California Fresno, California
Des Moines, Iowa Perth Amboy, New Jersey Orlando, Florida
Form 5639 K Product 2784 Made in U.S.A.
EPA Reg. No. 239-1296 AA

CELERY: Early and Late Blight—Use 30 to 50 lbs. per acre. Begin dusting when the plants first come through the ground in the seedbed. Make further applications at 3 to 5 day intervals. Continue applications after plants are set in the field on a regular weekly schedule. Remove residues by washing and stripping. Do not apply within 14 days of harvest.

CUCUMBERS, CANTALOUPE, WATERMELONS: Downy Mildew, Angular Leaf Spot, Anthracnose, Scab or Gummy Stem Blight—Use 30 lbs. per acre. Apply when disease first appears. Apply on weekly schedule or more often under severe disease conditions. Do not apply within 5 days of harvest.

HOPS: Downy Mildew—40 lbs. per acre. Dust on crowns when new growth starts. Begin vine dusting at first training and repeat at 5 to 10-day intervals. Do not apply within 14 days of harvest.

LETTUCE: Downy Mildew—40 to 50 lbs. per acre. Treat when disease first appears. Repeat as necessary. Avoid excessive applications shortly before harvest. Do not apply to lettuce within 10 days of harvest.

SPINACH: Downy Mildew—Use 30 lbs. per acre. Start applications as soon as plants emerge and repeat at 7 to 10-day intervals. Do not apply within 10 days of harvest.

STRAWBERRIES: Leaf Spot—40 to 50 lbs. per acre. Begin application before disease appears. Repeat at 7 to 10-day intervals. Do not apply within 7 days of harvest.

TOBACCO—Blue Mold in Plant Bed: When plants are the size of a dime or when the disease is first reported in the area, dust plants thoroughly every 3 or 4 days using 1½ pounds per 100 sq. yds. for 4 applications, followed by 4 to 8 applications using 2½ to 4 pounds per 100 sq. yds. of bed.

TOMATOES: Early and Late Blight—Use 30 to 50 lbs. per acre. Start dusting 30 days after first cluster bloom. Continue applications at 7 to 10-day intervals. If late blight threatens, earlier begin applications immediately. Do not apply within 5 days of harvest.

CONDITIONS OF SALE: 1. Chevron Chemical Company warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants and crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability except those assumed by Chevron under 1 above) resulting from handling, storage, and use of this material.

CAUTION

READ THE LABEL

KEEP OUT OF REACH OF CHILDREN

CAUTION: May Cause Irritation of Nose and Throat! Do not breathe dust. Do not store or transport with feed or food. **Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.**

Zineb 5 Dust



ORTHO

ACCEPTED
AUG 13 1973
FUNGICIDE AND REEDENTICIDE ACT
REGISTERED UNDER NO. 1972 SUBJECT
TO APPLICABLE COMMENTS

(FUNGICIDE)

Active Ingredient	By Wt.
Zineb*	5%
Inert Ingredients	95%
Zinc expressed as metallic	1.18%
Zinc ethylene bisdithiocarbamate	

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER.

BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE.

DIRECTIONS

This product is formulated for application by ground or air equipment.

BEANS: Rust, Anthracnose, Downy Mildew—Use 30 to 40 lbs. per acre. Start dusting immediately upon the appearance of the disease, or when the disease threatens and continue regularly applications at 7 to 10-day intervals. **Anthracnose:** Start dusting when the first blossoms appear and continue regular applications at 7 to 10 day intervals. Do not apply within 7 days of harvest.

BROCCOLI, CABBAGE: Downy Mildew—40 to 50 lbs. per acre. Begin applications before disease appears. Repeat at 7 to 10 day intervals. Do not apply within 7 days of harvest.

CARROTS: Late Blight—40 to 50 lbs. per acre. Apply at first signs of disease and repeat as necessary. Do not apply within 7 days of harvest. Carrots are to be used for food or feed.

NET WEIGHT 50 LBS.

Chevron Chemical Company

Orto Division, San Francisco, Calif. 94119
 Richmond, California Fresno, California
 Des Moines, Iowa North Amboy, New Jersey Orlando, Florida
 Farmington, N. C. Portland, Oregon Tulsa, Oklahoma
 Wichita, Kansas Dallas, Texas St. Louis, Missouri

CELERY: Early and Late Blight—Use 30 to 50 lbs. per acre. Begin dusting when the plants first come through the ground in the seedbed. Make further applications at 3 to 5 day intervals. Continue applications after plants are set in the field on a regular weekly schedule. Remove residues by washing and stripping. Do not apply within 14 days of harvest.

CUCUMBERS, CANTALOUPE, WATERMELONS: Downy Mildew, Angular Leaf Spot, Anthracnose, Scab or Gummy Stem Blight—Use 30 lbs. per acre. Apply when disease first appears. Apply on weekly schedule or more often under severe disease conditions. Do not apply within 5 days of harvest.

HOPS: Downy Mildew—40 lbs. per acre. Dust on crowns when new growth starts. Begin vine dusting at first training and repeat at 5 to 10-day intervals. Do not apply within 14 days of harvest.

LETTUCE: Downy Mildew—40 to 50 lbs. per acre. Treat when disease first appears. Repeat as necessary. Avoid late season applications shortly before harvest. Do not apply within 7 days of harvest.

SPINACH: Downy Mildew—Use 30 lbs. per acre. Start applications as soon as plants emerge and repeat at 7 to 10-day intervals. Do not apply within 10 days of harvest.


STRAWBERRIES: Leaf Spot—40 to 50 lbs. per acre. Begin application before disease appears. Repeat at 7 to 10-day intervals. Do not apply within 7 days of harvest.

TOBACCO—Blue Mold in Plant Bed: When plants are the size of a dime or when the disease is first reported in the area, dust plants thoroughly every 3 or 4 days using 1½ pounds per 100 sq. yds. for 4 applications, followed by 1 to 3 applications using 2½ to 4 pounds per 100 sq. yds. of bed.

TOMATOES: Early and Late Blight—Use 30 to 50 lbs. per acre. Start dusting 30 days after first planting. Continue applications at 7 to 10 day intervals. If late blight threatens, increase to 40 to 50 lbs. per acre. Do not apply within 7 days of harvest.

CONDITIONS OF SALE: This material is sold by the Company as a fungicide and is warranted to conform to the claims made on the label and is reasonably fit for uses specified thereon. The Company makes no warranty, express or implied, as to the efficacy, or the quality of the material. CHEVRON, ORTHO, a guarantee, or a representation, are not to be construed as a warranty of this material.

The material is sold by the Company and Chevron's control is limited to the material as sold. All risks in connection with the use of this material are assumed by the user, but are not limited to damage to plants and crops, which the material is applied, lack of complete control, and damage caused by drift to other plants and crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user assume all risks and liabilities. For more information, contact your local Chevron representative or Chevron Chemical Company, P.O. Box 1000, San Francisco, California 94119.



CAUTION

READ THE LABEL

KEEP OUT OF REACH OF CHILDREN

CAUTION: May Cause Irritation of Nose and Throat. Do not breathe dust. Do not store or transport with feed or food. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.