

# SEVIN<sup>\*</sup> Sulfur 5-40 Dust



# ORTHO

ACCEPTED  
MAY 18 1973  
UNDER THE FEDERAL INSECTICIDE  
AND FUNGICIDE ACT  
FOR ECONOMIC POISON REGISTER-  
ED UNDER NO. SUBJECT  
TO ATTACHED COMMENTS.  
237-1348

## (INSECTICIDE-FUNGICIDE)

Active Ingredients	By Wt.
Carbaryl (1-naphthyl N-methylcarbamate) .....	5%
Sulfur .....	40%
Inert Ingredients .....	55%
U. S. Pat. Nos. 2,903,478-3,009,855	

**READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.**

KEEP PESTICIDE IN ORIGINAL CONTAINER.

BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE.

This product is formulated for application by ground or air equipment.

**BEANS:** Mexican Bean Beetle, Powdery Mildew, Rust, Leafhoppers—20 pounds per acre. Lygus, Stink Bugs—40 pounds per acre. Apply thoroughly at first signs of insects or diseases. When necessary repeat every 7 to 10 days. May be used up to harvest.

**CORN:** Corn Earworm—30 to 40 lbs. per acre. Apply at 3-day intervals starting when silks first appear. Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee populations. Make 4 to 5 applications until silks dry or turn brown. May be applied up to harvest.

**GRAPES:** Grape Leaf Folder—1st Brood: 20 to 25 lbs. per acre just before first brood larvae emerge from the first rolls. 2nd and 3rd Broods: 20 to 40 pounds per acre. Grape

Leafhopper—20 pounds per acre. Thorough coverage is essential. May be applied up to harvest.

**PEANUTS:** Blister Beetles, Mexican Bean Beetle—10 to 20 lbs. per acre. Thrips, Leafhoppers, Three-cornered Alfalfa Hopper, Cucumber Beetles, Bean Leaf Beetle, Japanese Beetle, Alfalfa Caterpillar, Velvet Bean Caterpillar—20 lbs. per acre. Armyworm, Webworms, Corn Earworm, Stink Bugs—20 to 30 lbs. per acre. Grasshoppers—10 to 30 lbs. per acre. Rust—30 lbs. per acre. May be applied up to the day of harvest or grazing.



### READ THE LABEL

KEEP OUT OF REACH OF CHILDREN

**CAUTION:** HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN. Avoid Breathing Dust. Do not Take Internally. **Skin Contact May Be Harmful.** Avoid Contact. Wash hands and Face Before Eating. Take Shower or Bath After Work. Wear Regular Long Sleeved Work Clothing. Change to Clean Clothing Daily. **Note to Physician:** Carbaryl is a moderate, reversible, Cholinesterase inhibitor. Atropine is antidotal. Do not store or transport with feed or food. **AVOID CONTAMINATION OF FOOD, FEED, FEEDING TROUGHS AND WATERING RECEPTACLES.** Some fruits and other plants are susceptible to injury from sulfur under certain climatic conditions. The user is advised not to use sulfur on any crop unless local use has proved that sulfur does not damage crops in that locality. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

For protection of honeybees avoid unnecessary use during periods when honeybees are visiting the crop. When necessary to use during such periods, warn bee-keepers well in advance to locate hives at a safe distance until one week after application.

**PLUMS, PRUNES:** Fruit Tree Leaf Roller, Peach Twig Borer, Plum Curculio, Red-Banded Leaf Roller, Codling Moth, Eastern Tent Caterpillar, Orange Tortrix, Western Tussock Moth—60 to 70 lbs. per acre. Apply when insects appear and repeat as necessary. Make full coverage application. For Codling Moth and Plum Curculio—Apply at petal fall and every 10 to 14 days thereafter until control is achieved. Do not apply within 1 day of harvest.

**POTATOES:** Colorado Potato Beetle—10 pounds per acre. Leafhopper and Flea Beetle—20 pounds per acre. Lygus Bugs—40 pounds per acre. Apply when insects appear. Apply dust thoroughly to upper and lower leaf surfaces. If needed, repeat applications at 7 to 10-day intervals.

**TOMATOES:** Tomato Fruitworm, Fall Armyworm, Flea Beetles, Russet Mite, Tomato Hornworm, European Corn Borer, Stink Bugs—50 to 60 lbs. per acre. Apply thoroughly at first signs of insect infestation. Repeat as necessary. May be applied up to the day of harvest. Because of the sulfur content of this dust, first obtain permission from your canner or processor before using it on tomatoes intended for sale to such canner or processor.

**CONDITIONS OF SALE:** 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Chevron under 1 above) resulting from handling, storage, and use of this material.

\*SEVIN trademark of Union Carbide Corporation for the active ingredient, 1-naphthyl N-methylcarbamate.

## NET WEIGHT 50 LBS.

**Chevron Chemical Company**

Ortho Division/San Francisco, Calif. 94119

Richmond, California

Fresno, California

Des Moines, Iowa

Perth Amboy, New Jersey

Orlando, Florida

Form 5712-Z

Product 2831

Made in U.S.A.

EPA Reg. No. 239-1348-AA