


THIODAN Sulfur 3-50 Dust



ORTHO

(INSECTICIDE-FUNGICIDE)

Active Ingredients	By Wt.
*Endosulfan	3%
Sulfur	50%
Inert Ingredients	47%
*Hexachlorohexahydromethano-2,4,3-benzodioxathiepin oxide	
Sulfur Fineness Through 325 Mesh	93% Typical
U. S. Pat. No. 2,799,685	

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER.

BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE.

DIRECTIONS

Not for use or storage in or around the home.

This product is formulated for application by ground or air equipment.

ALFALFA SEED CROPS: Spittlebug, Aphids, Lygus Bugs, Mites—50 lbs. per acre.

Crop to be used for seed production only. Do not feed treated forage.

COTTON: Lygus Bugs, Flea Beetles, Aphids, Leafhoppers, Stink Bugs—25 to 30 pounds per acre. Apply as insects appear and repeat as necessary. Thorough coverage is essential. Do not apply after bolls open. Do not graze meat or dairy animals in treated fields. Workers entering treated areas within 24 hours of treatment should wear

protective clothing. Do not plant root crops other than melons, potatoes, sweet potatoes and sugarcane in follow-up crops.

GRAPES: Grape Leafhopper, Powdery Mildew—35 to 45 lbs. per acre. Apply when insects or disease first appear and repeat as necessary. Do not use on Concord variety of Grapes as severe injury is likely to occur. Do not apply within 7 days of harvest.



READ THE LABEL

KEEP OUT OF REACH OF CHILDREN

CAUTION: Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing dust. Avoid contact with skin, eyes or clothing. In case of contact with skin or eyes, flush with plenty of water; for eyes, get medical attention. Wash with soap and water after handling and before eating or smoking; wear clean clothing. Do not store or transport near feed or food. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals. Workers entering treated areas the day of application should wear protective clothing. Use of this material in a manner or at a time other than in accordance with the directions may produce plant injury. Some fruits and other plants are susceptible to injury from sulfur under certain climatic conditions. The user is advised not to use sulfur on any crop unless local use has proved that sulfur does not damage crops in that locality. **Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.**

Highly toxic to fish and wildlife. Birds feeding on treated areas may be killed. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is toxic to bees and should not be applied if bees are actively visiting the area.

ORNAMENTALS, FLOWER SEED CROPS: Aphids (except Cabbage Aphid and Woolly Apple Aphid), Powdery Mildew—25 to 30 pounds per acre. Apply when aphids or disease appear and repeat as necessary. Thorough coverage is essential.

PEACHES: Green Peach Aphid, Stink Bugs - 50 lbs. per acre. Do not apply within 30 days of harvest nor make more than 2 applications during fruiting period.

POTATOES: Aphids, Potato Flea Beetle, Potato Leafhopper, Colorado Potato Beetle—Apply 30 to 35 pounds per acre. Apply when insects first appear and repeat as necessary. Thorough coverage is essential. Do not plant root crops other than carrots, potatoes, sweet potatoes and sugar beets as follow-up crops.

STRAWBERRIES: Cyclamen Mite - 60 to 70 pounds per acre. Apply as a crown treatment at the pre-bloom stage in the spring before the formation of buds. Do not re-apply within 35 days when fruit is present. Do not apply within 4 days of harvest. Dust crown of each plant thoroughly. Repeat treatment after final harvest if mites are present.

CONDITIONS OF SALE: 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Chevron makes no warranty, express or implied, that the material is suitable for use in any particular situation, with the exception of those specifically stated on the label. Users should consult the label for directions to which they should refer for proper use. Chevron's liability for any loss or damage, including consequential damages, shall be limited to the amount of the purchase price of the material. Such loss or damage shall be limited to the amount of the purchase price of the material. Users should consult the label for directions to which they should refer for proper use. Buyers should consult the label for directions to which they should refer for proper use. Chevron's liability for any loss or damage, including consequential damages, shall be limited to the amount of the purchase price of the material.

NET WEIGHT 50 LBS.

Chevron Chemical Company

Ortho Division San Francisco, Calif. 94119

Richmond, California Fresno, California Honolulu, Hawaii

Des Moines, Iowa Perth Amboy, New Jersey Orlando, Florida

Form 5/71 M EPA Reg. No. 239.1354 Product 2830 Made in U.S.A.