



SEVIN\*

Sulfur 7.5-50 Dust

MAY 6, 1971

[INSECTICIDE-FUNGICIDE]
NET WEIGHT 50 LBS.

Table with 2 columns: Active Ingredients, Inert Ingredients, and By Wt. Percentages.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP OUT OF REACH OF CHILDREN

REPEAT APPLICATIONS WHEN NECESSARY. REPEAT APPLICATIONS SHOULD BE MADE AT 7 TO 10 DAY INTERVALS.

DIRECTIONS

For general use, dilute 1 part of product with 100 parts of water in a spray or mist.

BEANS: Mexican Bean Beetle—10 to 20 lbs. per acre. Leafhoppers—15 lbs. per acre. Lygus, Stink Bugs, Powdery Mildew—20 lbs. per acre. Apply when insects appear and repeat at 7 to 10 day intervals. When necessary repeat every 7 to 10 days. May be used up to harvest.

CORN: Corn Earworm—10 to 20 lbs. per acre. Apply at 3-day intervals starting when silks first appear. Application of this product to the tassel region of corn plants in a pre-silking period will effectively reduce bee populations. Make 2 to 3 applications in the early or tassel stage. May be applied up to harvest.

COTTON: Early Season—Thrips, Cotton Fleahopper—10 to 15 lbs. per acre. Apply when insects appear. Mid and Late Season—Boll Weevil, Bollworm, Pink Bollworm, Cotton Leafworm, Lygus Bugs, Cotton Leaf Perforator—10 to 25 lbs. per acre. Apply when insects appear and repeat at 7 to 10 day intervals. May be used up to harvest. Do not apply with this insecticide before or during the schedule of treatments.



READ THE LABEL

KEEP OUT OF REACH OF CHILDREN

CAUTION: HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN. Avoid Breathing Dust. Do Not Take Internally. Skin Contact May Be Harmful. Avoid Contact. Wash Hands and Face Before Eating. Take Shower or Bath After Work. Wear Protective Clothing. Note for Physician: In case of moderate to severe poisoning, treatment should be symptomatic and supportive.

Do not use in areas where the dust may be inhaled. Do not use in areas where the dust may be inhaled. Do not use in areas where the dust may be inhaled.

Do not use in areas where the dust may be inhaled. Do not use in areas where the dust may be inhaled. Do not use in areas where the dust may be inhaled.

GRAPES: Grape Leaf Fleahopper—1st Brood—10 to 15 lbs. per acre. Apply when insects appear. 2nd and 3rd Broods—15 lbs. per acre. Apply when insects appear. Leafhopper, Powdery Mildew—20 lbs. per acre. Apply when insects appear and repeat at 7 to 10 day intervals. May be used up to harvest.

PEACHES: Oriental Fruit Moth, Peach Twig Borer, Plum Curculio, Cat-Facing Insects, Periodical Cicada, Powdery Mildew—20 lbs. per acre. Apply when insects appear and repeat at 7 to 10 day intervals. May be used up to harvest.

POTATOES: Colorado Potato Beetle—10 lbs. per acre. Leafhopper and Flea Beetle—15 lbs. per acre. Lygus Bugs—25 lbs. per acre. Apply when insects appear. Apply dust thoroughly to upper and lower leaf surfaces. Repeat application at 7 to 10 day intervals.

TOMATOES: Tomato Fruitworm, Fleahopper, Flea Beetle, Russet Mite, Tomato Plantworm, European Corn Borer, Stink Bug—20 lbs. per acre. Apply when insects appear and repeat at 7 to 10 day intervals. May be used up to harvest.

CONDITIONS OF SALE: This product is sold as a general use insecticide and fungicide. It is not intended for use as a fumigant or for control of insects in stored products. It is not intended for use on ornamental plants or in greenhouses.

Do not use in areas where the dust may be inhaled. Do not use in areas where the dust may be inhaled. Do not use in areas where the dust may be inhaled.

Do not use in areas where the dust may be inhaled. Do not use in areas where the dust may be inhaled. Do not use in areas where the dust may be inhaled.

CHEVRON CHEMICAL COMPANY

FORM 5827-7

USDA Reg. No. 239-1360

ORTHO DIVISION

San Francisco, California 94119

FORM 5827-7

FORM 5827-7

RICHMOND, CALIFORNIA

PORTLAND, OREGON

SAN FRANCISCO, CALIFORNIA



DES MOINES, IOWA

CHERRY HILL, NEW JERSEY