


KELTHANE® 4 Dust



ORTHO



(MITICIDE)

Active Ingredient By Wt.
1,1-bis(p-chlorophenyl)-2,2,2-trichloroethanol 4%
Inert Ingredients 96%

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER.

BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE.

This product is formulated for application by ground or air equipment.

APPLES: Spider Mites—60 to 70 lbs. per acre. Apply at petal fall to second cover or whenever mite population appears. If repeat application is necessary, wait 30 days. Do not apply within 7 days of harvest.

BEANS: Spider Mites—30 to 37 lbs. per acre by ground or air application. Do not feed treated vines to meat or dairy animals. Do not apply within 7 days of harvest.

CANEBERRIES (Blackberries, Boysenberries, Dewberries, Loganberries, Raspberries): Two-Spotted Mite, Pacific Mite—30 to 40 lbs. per acre. Apply at petal fall to second cover or whenever mite population appears. Do not apply within 7 days of harvest.

CANTALOUPEs, CUCUMBERS, SQUASH, MELONS, WATER-MELONS: Two-Spotted Mite, Pacific Mite—35 lbs. per acre.

NET WEIGHT 50 LBS.

Chevron Chemical Company

Ortho Division, San Francisco, Calif. 94119

Richmond, California Fresno, California Honolulu, Hawaii
Des Moines, Iowa Perth Amboy, New Jersey Orlando, Florida

Form 5291-C Product 2880 Made in U.S.A.

EPA Reg. No. 239-1389-AA

Apply at first sign of mites and repeat as necessary. Do not apply within 2 days of harvest.

CITRUS: Citrus Red Mite, Citrus Flat Mite, Six Spotted Mite, Yuma Mite, Two-Spotted Mites, Pacific Mite—50 to 70 lbs. per acre. Apply when mites first appear and repeat as necessary. Do not apply within 7 days of harvest.

COTTON: Pacific Mite, Atlantic Mite, Two-Spotted Mite, Red Spider Mites—40 lbs. per acre. Do not apply within 14 days of harvest. Do not graze treated fields by meat or dairy animals.

FIGS, PLUMS, PRUNES: Spider Mites—35 to 50 lbs. per acre. Apply at petal fall to first cover or whenever mite population appears. Do not apply within 7 days of harvest. Do not repeat application within 30 days.

GRAPES: Spider Mites—35 lbs. per acre. Apply as mites appear and repeat as necessary. Do not apply within 7 days of harvest.

HOPS: Two-spotted and Pacific Mite—35 lbs. per acre. Apply when first mites appear, repeat as necessary. Do not apply within 7 days of harvest.

PEACHES, NECTARINES: Spider Mites—50 to 60 lbs. per acre. Apply at petal fall to first cover or whenever mite population appears. Do not apply within 7 days of harvest.



READ THE LABEL

KEEP OUT OF REACH OF CHILDREN

CAUTION: Avoid breathing dust. Do not use in or near occupied buildings. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crop, meadow rendered unfit for sale, use or consumption.

This product is toxic to fish. Keep out of any body of water. Do not use in or near occupied buildings. Do not use in or near occupied buildings.

PEARS: European Red Mites, Two-Spotted Mites, Brown Almond Mites, Pacific Mites—45 to 50 lbs. per acre. Apply at petal fall to second cover or whenever mite population appears. Do not apply within 7 days of harvest.

SEED ALFALFA: Two-spotted and Pacific Mite — Ground Application Airplane Application
Apply at petal fall to first cover or whenever mite population appears. Repeat application as necessary. Do not apply within 7 days of harvest. Do not graze treated fields by meat or dairy animals. Do not apply within 7 days of harvest.

STRAWBERRIES: Two-Spotted Mite, Pacific Mite—50 lbs. per acre. Apply when mites appear and repeat as necessary. Do not apply within 2 days of harvest.

WALNUTS: European Red Mites, Two-Spotted Mites and Brown Almond Mites—40 to 50 lbs. per acre. Apply as mites appear and repeat as necessary. Thorough coverage is essential. Do not apply within 4 days of harvest.

CONDITIONS OF SALE: This product is sold by the Company as a commodity and is warranted to conform to the chemical description on the label and is reasonably fit for use as directed thereon. The Company makes, nor authorizes any agent or representative to make, any other warranty of performance, MERCHANTABILITY, guarantee or representation in connection with the sale of this material.

Damage caused by use of this material, beyond Chevron's control, is sold as is, with all risks in connection with its use. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, damage to property, and damage to the environment. Such risks occur even though the material is used in accordance with the directions stated hereon and the user has followed the directions and used the material in accordance with the directions. The user assumes all risks and liability for damage caused by the use of this material under the conditions stated above. The user assumes all liability for damage caused by the use of this material.