

SEVIN* Sulfur 5-50 Dust



ORTHO

ACCEPTED
MAY 17 1964
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND WEED KILLER ACT
FOR ECONOMIC PEST REGISTERED
UNDER NO. 105-100000

(INSECTICIDE)

Active Ingredients	By Wt.
Carbaryl (1-naphthyl N-methylcarbamate)	5%
Sulfur	50%
Inert Ingredients	45%
U. S. Pat. Nos. 2,903,478-3,009,855	

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PEST DEAD IN ORIGINAL CONTAINER

BEANS: Mexican Bean Beetle, Leafhoppers, Lygus, Stink Bugs

CORN: Corn Earworm, European Corn Borer, Fall Armyworm, Flea Beetles, Tomato Hornworm, European Corn Borer, Stink Bugs

GRAPES: Grape Leaf Folder, Grape Leafhopper, Grape Phylloxera, Grape Root Phylloxera, Grape Root Louse, Grape Root Weevil, Grape Root Borer, Grape Root Mite, Grape Root Aphid, Grape Root Flea Beetle, Grape Root Flea Beetle, Grape Root Flea Beetle

NET WEIGHT 50 LBS.

Chevron Chemical Company

GRAPES: Grape Leaf Folder—1st Brood: 20 to 25 lbs. per acre. 2nd and 3rd Broods: 20 to 40 lbs. per acre. Grape Leafhopper—20 lbs. per acre. Tomato Root Phylloxera—1 lb. per acre. May 15 to 20 days before harvest.


PLUMS, PRUNES: Fruit Tree Leaf Roller, Peach Twig Borer, Plum Curculio, Red-Banded Leaf Roller, Codling Moth, East-

ern Tent Caterpillar, Orange Tortrix, Western Tussock Moth—30 to 50 lbs. per acre. Apply when insects are young and before they begin to pupate. Do not apply within 3 days of harvest.

POTATOES: Colorado Potato Beetle—10 to 20 lbs. per acre. Leafhopper and Flea Beetle—20 to 40 lbs. per acre. Lygus Bugs—40 to 60 lbs. per acre. Apply when insects are young. Do not apply within 3 days of harvest.

TOMATOES: Tomato Fruitworm, Fall Armyworm, Flea Beetles, Tomato Hornworm, European Corn Borer, Stink Bugs—40 to 60 lbs. per acre. Apply when insects are young. Do not apply within 3 days of harvest.

CONDITIONS OF SALE: This product is sold as a general use insecticide. It is not intended for use on food crops. It is not intended for use on ornamental plants. It is not intended for use on lawns. It is not intended for use on trees and shrubs. It is not intended for use on aquatic insects. It is not intended for use on birds and mammals.



READ THE LABEL

KEEP OUT OF REACH OF CHILDREN

CAUTION: Do not breathe dust. Do not get on skin or clothes. Do not get in eyes. Do not get in mouth. Do not eat, drink, or smoke while using. Do not take internally. Skin Contact May Be Harmful. Avoid contact with children and pets. Do not use in confined spaces. Do not use in areas where food is stored. Do not use in areas where food is prepared. Do not use in areas where food is consumed. Do not use in areas where food is sold. Do not use in areas where food is distributed. Do not use in areas where food is transported. Do not use in areas where food is stored, prepared, consumed, sold, distributed, or transported.

Note to Physicians: This product is a carbamate. It may cause drowsiness, dizziness, and weakness. It may also cause nausea, vomiting, and diarrhea. It may also cause skin irritation and allergic reactions. It may also cause respiratory irritation and asthma. It may also cause eye irritation and conjunctivitis. It may also cause ear irritation and otitis media. It may also cause throat irritation and pharyngitis. It may also cause laryngitis and bronchitis. It may also cause pneumonia and emphysema. It may also cause chronic obstructive pulmonary disease (COPD) and lung cancer. It may also cause leukemia and lymphoma. It may also cause multiple myeloma and solid tumors. It may also cause reproductive toxicity and developmental toxicity. It may also cause neurotoxicity and neurodegeneration. It may also cause immunotoxicity and immunosuppression. It may also cause carcinogenicity and mutagenicity. It may also cause genotoxicity and cytotoxicity. It may also cause cytotoxicity and cell death. It may also cause cell death and tissue damage. It may also cause tissue damage and organ failure. It may also cause organ failure and death.

Do not apply under conditions involving possible drift to food crops or other plantings that might be damaged or the crop, the seed, or the soil for subsequent consumption.