

FACSIMILE OF EXACT COPY ON LATEST BAG PRINTING.

ACCEPTED
SEP 8 1969
UNDER THE FEDERAL INSECTICIDE
FUNCTIONAL AND RESISTANCE ACT
FOR ECONOMIC POISON REGISTER-
ING UNDER NO. 239-1470



SEVIN* Bean Dust

[INSECTICIDE]

Active Ingredient
Carbaryl (1-naphthyl N-methylcarbamate)
Inert Ingredients
U. S. Pat. Nos. 2,903,478-3,009,855

By Wt.
1.75%
98.25%



READ THE LABEL

KEEP OUT OF REACH OF CHILDREN

CAUTION: HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN. Avoid Breathing Dust. Do Not Take Internally. Skin Contact May Be Harmful. Avoid Contact. Wash Hands and Face Before Eating. Take Shower or Bath After Work. Wear Regular Long Sleeved Work Clothing. Change to Clean Clothing Daily. Note for Physician: Carbaryl is a moderate reversible, Cholinesterase inhibitor. Atropine is antidotal. **AVOID CONTAMINATION OF FOOD, FEED, FEEDING TROUGHS AND WATERING RECEPTACLES.** This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. Do not apply under conditions involving possible harm to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

*Trade mark of Union Carbide Corporation for the active ingredient,
1-naphthyl N-methylcarbamate

**NET WEIGHT
4 POUNDS**

DIRECTIONS ON BACK

**BURN BAG IMMEDIATELY
WHEN EMPTY.
STAY OUT OF SMOKE.**

FORM 1414H (5-6-69) (S.A. Reg. No. 141570) (E.P. Reg. No. 124) Made in U.S.A.
CHEVRON CHEMICAL COMPANY
ORTHO DIVISION - P.O. Box 1170, Richmond, Va. 23261

FACSIMILE OF EXACT COPY ON LATEST BAG PRINTING

ORTHO[®]
SEVIN* BIOCIN DUST

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER.

DIRECTIONS

BEANS: Mexican Bean Beetle — $\frac{3}{8}$ lb. per 1,000 sq. ft. Leafhoppers, Bean Leaf Beetle, Cucumber Beetles, Western Bean Cutworm, Velvetbean Caterpillar, Flea Beetles, Japanese Beetle — $\frac{1}{4}$ lbs. per 1,000 sq. ft. Corn Earworm, Armyworms, Tarnished Plant Bug, Stink Bugs — $\frac{1}{4}$ to $\frac{1}{2}$ lbs. per 1,000 sq. ft. May be used up to harvest. Apply thoroughly at first signs of infestation. When necessary repeat every 7 to 10 days.

POTATOES: Colorado Potato Beetle — $\frac{3}{8}$ lb. per 1,000 sq. ft. Leafhopper and Flea Beetle — $\frac{1}{4}$ lbs. per 1,000 sq. ft. Apply when insects appear. Apply dust thoroughly to upper and lower leaf surfaces. If needed, repeat applications at 7 to 10-day intervals.

CONDITIONS OF SALE: 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated thereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Chevron under 1 above) resulting from handling, storage, and use of this material.

FORM 6114-H