

SEVIN* Sulfur 10-50 Dust



ORTHO

ACCEPTED

AUG 24 1973

UNDER THE FEDERAL LAW FOR
FUNGICIDE AND PESTICIDE
FOR ECONOMIC PEST CONTROL
ED UNDER NO. 100-100000

(INSECTICIDE-FUNGICIDE)

Active Ingredients	By Wt.
Carbaryl (1-naphthyl N-methylcarbamate)	10%
Sulfur	50%
Inert Ingredients	40%
Sulfur Fineness Through 325 Mesh.....	93% Typical
U. S. Pat. Nos. 2,903,478-3,009,855	

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER.

BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE.

This product is formulated for application by ground or air equipment.

BEANS: Mexican Bean Beetle, Leafhoppers, Powdery Mildew—10 pounds per acre. Lygus, Stink Bugs—20 pounds per acre. Apply thoroughly at first signs of disease or insect infestation. When necessary repeat every 7 to 10 days. May be used up to harvest.

CORN: Corn Earworm—15 to 20 lbs. per acre. Apply at 7-day intervals starting when silks first appear. Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee populations. Make application until silks dry or turn brown. May be applied up to harvest.

NET WEIGHT 50 LBS.

Chevron Chemical Company

Ortho Division/San Francisco, California 94119
Richmond, California Fresno, California
Des Moines, Iowa Perth Amboy, New Jersey Orlando, Florida
Form 6265-W Product 3153 Made in U.S.A.

EPA Reg. No. 239-1601-AA

COTTON: Early-Season—Thrips, Cotton Fleahopper—8 to 15 lbs. per acre as soon as insects appear. Mid and Late-Season—Boll Weevil, Bollworm, Pink Bollworm, Cotton Leafworm, Lygus Bugs, Cotton Leaf Perforator—20 lbs. per acre. Apply when insects appear and repeat at 5 to 7-day intervals. Aphid populations will usually be suppressed when this insecticide is used on a regular schedule of treatments.

GRAPES: Grape Leaf Folder—1st Brood: 10 to 15 lbs. per acre just before first brood larvae emerge from the first rolls. 2nd and 3rd Brood: 10 to 20 lbs. per acre. Grape Leafhopper, Powdery Mildew—10 lbs. per acre. Apply thoroughly at first signs of disease or insect infestation. When necessary repeat

every 7 to 10 days. Thorough coverage is essential. May be applied up to harvest.

PEACHES: Oriental Fruit Moth, Peach Twig Borer, Plum Curculio, Cat-Facing Insects, Periodical Cicada, Powdery Mildew, Brown Rot—35 to 40 lbs. per acre. Apply thoroughly at first signs of disease or insect infestation. When necessary repeat every 7 to 10 days. Do not apply within 1 day of harvest.

PLUMS, PRUNES: Fruit Tree Leaf Roller, Peach Twig Borer, Plum Curculio, Red-Banded Leaf Roller, Codling Moth, Eastern Tent Caterpillar, Orange Tortrix, Western Tussock Moth—40 to 50 lbs. per acre. Apply when insects appear and repeat as necessary. Make full coverage application. Do not apply within 1 day of harvest.


POTATOES: Colorado Potato Beetle—10 lbs. per acre. Leafhopper, Flea Beetle, Lygus Bugs—20 lbs. per acre. Apply when insects appear. Apply dust thoroughly to upper and lower leaf surfaces. If needed, repeat applications at 7 to 10-day intervals.

TOMATOES: Tomato Fruitworm, Fall Armyworm, Flea Beetles, Russet Mite, Tomato Hornworm, European Corn Borer, Stink Bugs—30 to 40 lbs. per acre. Apply thoroughly at first signs of insect infestation. Repeat as necessary. May be applied up to the day of harvest. Because of the sulfur content of this dust, first obtain permission from your canner or processor before using it on tomatoes intended for sale to such canner or processor.

CONDITIONS OF SALE: 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Chevron under 1 above) resulting from handling, storage, and use of this material.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.



READ THE LABEL

KEEP OUT OF REACH OF CHILDREN

CAUTION: HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN. Avoid Breathing of Dust. Do Not Take Internally. Skin Contact May Be Harmful. Avoid Contact. Wash Hands and Face Before Eating. Take Shower or Bath After Work. Wear Regular Long Sleeved Work Clothing. Change to Clean Clothing Daily.

Note for Physician: Carbaryl is a moderately potent insecticide and neurotoxin. Atropine is antidotal.

Do not use on or transport near food or feed. AVOID CONTACT WITH INGESTIBLE FOOD, FEED, FEEDING TROUGHS AND WATERING BELLS. Stink bugs and other insects may be susceptible to injury from water and other materials. This product has been used not to cause injury on any crop unless, however, it has been proved that sulfur does not damage crops in that locale. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.