

ORTHO[®] Sulfur 15-25 Dust



ORTHO

ACCEPTED
Sept 4, 1973
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR FOLIAR MITICIDE REGISTER-
ED UNDER NO. 239-1668

(FUNGICIDE)

Active Ingredients	By Wt.
Captan*	15%
Sulfur	25%
Inert Ingredients	60%
* N-[(trichloromethyl)thio]-4-cyclohexene-1,2-dicarboximide.	

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER.

STORE IN COOL, DRY PLACE. PROTECT FROM EXCESSIVE HEAT.

DO NOT REUSE EMPTY CONTAINER. DESTROY IT BY BURYING IT IN WASTE OR BURNING. STAY AWAY FROM SMOKE OR FUMES.

This product is formulated for application by ground or air equipment.

GRAPES: Botrytis Rot (Bunch Rot), Powdery Mildew, Silver Mite, Storage Rots—33 pounds per acre. Make 2 applications before bloom and 1 application just after bloom. Repeat periodically making up to 3 cover applications before bunches close.

CELERY: Pink Rot, Damp-Off—(Field)—13 lbs. per acre. (Seed Bed)—25 to 30 pounds per acre. Apply

NET WEIGHT 50 LBS.

Chevron Chemical Company

Ortho Division/San Francisco, Calif. 94119
Richmond, California Fresno, California
Des Moines, Iowa Perth Amboy, New Jersey Orlando, Florida
Form 6365-K Product 3142 Made in U.S.A.
EPA Reg. No. 239-1668-AA

when all seeds have germinated. Repeat as necessary.

PEACHES: Brown Rot, Storage Rots, Powdery Mildew, Silver Mite, Peach Scab—33 pounds per acre. Make 2 to 3 prebloom applications. Repeat at petal fall and 3 to 4 covers as necessary. Make preharvest application where necessary.

POTATOES: Early and Late Blight—Begin applications when plants are 2 to 4 inches high. Apply 15 to 20 pounds per acre.

ROSES (Field): Rust, Powdery Mildew—35 to 40 pounds per acre. Apply to give even coverage of upper and lower leaf surfaces at first sign of disease. Repeat at regular intervals as necessary to maintain control.



READ THE LABEL

KEEP OUT OF REACH OF CHILDREN

CAUTION: Avoid inhalation of dust. Avoid contact with skin. Do not store or transport near feed or food. Some fruits and other plants are susceptible to injury from sulfur under certain climatic conditions. The user is advised not to use sulfur on any crop unless local use has proved that sulfur does not damage crops in that locality. **Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.**

This product is toxic to fish. Keep out of lakes, streams or ponds.

STRAWBERRIES: Botrytis Rot, Powdery Mildew—20 pounds per acre immediately after picking. Repeat applications after picking as necessary.

TOMATOES (Established Plants): Early Blight, Late Blight, Anthracnose, Russet Mite—On young plants, apply 15 to 20 lbs. per acre, increasing to 25 lbs. per acre on mature plants. Apply at first sign of insects and diseases. Because of the sulfur content of this dust, first obtain permission from your canner or processor before using it on tomatoes intended for sale to such canner or processor.

COMPATIBILITY: Do not use this product on foliage closely following oil sprays.

CONDITIONS OF SALE: 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Chevron under 1 above) resulting from handling, storage, and use of this material.