

KELTHANE[®]-Sulfur 4-30 Dust



ORTHO

(MITICIDE-FUNGICIDE)

Active Ingredients	By Wt.
1,1-bis (p-chlorophenyl)-2,2,2-trichloroethanol	4%
Sulfur	30%
Inert Ingredients	66%

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER.

BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE.

DIRECTIONS

This product is formulated for application by ground or air equipment.

ALFALFA (Seed): Two-Spotted Mite, Strawberry (Atlantic) Mite, Carmine Mite, Pacific Mite—25 to 35 lbs. per acre. Use restricted to California, Idaho, Oregon, Nevada and Washington. Fields not to be grazed by livestock. Hay containing residues not to be fed or baled. The field must be plowed after harvest, by burning or by deep plowing.

APPLES: Spider Mites—60 to 70 lbs. per acre. Apply at petal fall to second cover or whenever mite population appears. If repeat application is necessary, wait 30 days. Do not apply within 7 days of harvest.

NET WEIGHT 50 LBS.


Chevron Chemical Company

Ortho Division, San Francisco, Calif. 94119

Richmond, California Fresno, California Honolulu, Hawaii
Des Moines, Iowa Perth Amboy, New Jersey Orlando, Florida

Form 6295-N Product 3164 Made in U.S.A.

© A Reg. No. 259-1600-AA



BEANS: Spider Mites—30 to 37 lbs. per acre by ground or air application. Do not feed treated vines to meat or dairy animals. Do not apply within 7 days of harvest.

FIGS: Red Spider Mites—50 pounds per acre. Apply at petal fall to first cover or whenever mite population appears. If repeat application is necessary, wait 30 days. Do not apply within 7 days of harvest.

GRAPES: Powdery Mildew, Spider Mites—35 lbs. per acre. Apply at first sign of insects or diseases. Repeat at 10 to

14 day intervals as necessary. Do not apply within 7 days of harvest.

HOPS: Two-Spotted and Pacific Mite—35 lbs. per acre. Apply when first mites appear. Repeat as necessary. Do not apply within 7 days of harvest.

STRAWBERRIES: Red Spider Mites—40 to 50 pounds per acre. Apply when mites appear and repeat as necessary. Do not apply within 7 days of harvest.

CONDITIONS OF SALE: Chevron Chemical Company ("Chevron") warrants that the material herein conforms to the chemical description on the label and is reasonably fit for use as directed, to the extent that such use is made in accordance with the label directions. Chevron makes no other warranty, expressed or implied, and does not warrant the merchantability of the material.

Critical and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, risk of incomplete control and damage caused by drift to other plants or crops. Such risks are known even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Chevron under 1 above) resulting from handling, storage, and use of this material.

CAUTION **READ THE LABEL**

KEEP OUT OF REACH OF CHILDREN

CAUTION: Avoid breathing dust. Do not store or transport near feed or food. Some fruits and other plants are susceptible to injury from sulfur under certain climatic conditions. The user is advised not to use sulfur on any crop unless local use has proved that sulfur does not damage crops in that locality. **Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.**

Highly toxic to fish. Do not contaminate any body of water nor apply to any area not specified on the label.