




ORTHO SEVIN* Garden Spray



ORTHO

Makes Up To
15 Gals. Diluted
Spray

SEVIN* Garden Spray

READ ENTIRE LABEL USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS AND DIRECTIONS
KEEP PESTICIDE IN ORIGINAL CONTAINER DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK
CONTAINERS

DIRECTIONS Use 2 level Tablespoonfuls ORTHO SEVIN Garden Spray in 1 gallon of water. For best results apply with an ORTHO SPRAYETTE. To use mix designated amount of ORTHO SEVIN Garden Spray in a little water to make a thin paste, then add required amount of water. Do not store diluted spray. Make new dilution for each use.

ROSES, ORNAMENTALS, FLOWERS, SHRUBS, SHADE TREES Japanese Beetles, Aphids, Bagworm, Gypsy Moth, Elm Leaf Beetle, Certain Leafminers, Rose Slug, Periodical Cicada (17-year locust), Scale Insects, Tent Caterpillars, Blister Beetles, Leaf Rollers, Exposed Thrips—Start spraying when insects or their damage appear. Repeat weekly or as necessary. Do not use on Boston Ivy.

FRUIT TREES (APPLES, PEARS AND PEACHES) Apple Maggot, Plum Curculio, Apple Aphid, Leaf Rollers, Peach Tree Borer, Oriental Fruit Moth, Codling Moth—Make one application between bud break and bloom, a second application at petal fall, and then at 3 to 4 week intervals for 3 to 4 months as necessary. **Apple Maggot**—For best control, apply at 10 day intervals. **Rosy Apple Aphid**—Apply before leaves are curled. **NOTE:** To avoid foliage injury on McIntosh apples and possible fruit thinning on all varieties of apples, delay use until at least 30 days after bloom. Do not apply within 1 day of harvest.

VEGETABLES (TOMATOES, EGGPLANT, PEPPERS, CUCUMBERS, SUMMER SQUASH) Pickleworm, Fruit worms, Melon Worms, Hornworms, Fall Armyworms, Corn Borer—Apply when insects first appear. Repeat at 7 to 10 day intervals. For Corn Borer control apply thoroughly into the whorls when plant is about knee high. Repeat at weekly intervals until control is achieved.

LAWN INSECTS Sod Webworms, Fall Armyworms, Earwigs, Grasshoppers, Fleas and Ants—Use 3 ounces (18 level Tablespoonfuls) in 15 gals. water and cover 500 sq. ft. of established lawn. For best results, use an ORTHO Lawn Sprayer. Mow lawn first and apply after rain or watering. Do not water for at least two days after application. Repeat in 2 or 3 weeks or as necessary.

CAUTION HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN. Avoid Breathing Dust or Spray. Avoid Skin Contact. Avoid Eye Contact. In case of eye contact, immediately flush eyes with plenty of fresh water. Wash Hands and Face Before Eating. Keep children and animals off treated lawns until these areas are dry.

Note to Physicians: Emergency Information—call (415) 233-3737. Carbaryl is a moderate, reversible, Cholinesterase inhibitor. Atropine is antidotal.

AVOID CONTAMINATION OF FOOD. Dispose of container promptly when empty.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

NOTICE Buyer assumes all responsibility for safety and use not in accordance with directions.

Note: This product is sold by weight and package is full when packed, but due to fluffy nature the material is inclined to settle.

SEVIN* Trade Mark of Union Carbide Corporation for the active ingredient: 1-naphthyl N-methylcarbamate

Chevron Chemical Company

Ortho Division, San Francisco, CA 94119 Richmond, CA 94804

Product 3333 Made in U.S.A.

Form 6503-S

EPA Est. 239-MU-1

EPA Reg. No. 239-1701AA



Active Ingredient
Carbaryl (1-naphthyl N-methylcarbamate) 50%
Inert Ingredients 50%
U.S. Pat. No. 2,933,478; 3,009,855

CAUTION: Keep out of reach of children.
See back panel for additional cautions

NET WT. 6 OZ.