



DIFOLATAN[®] 80 Wettable

NET WEIGHT 50 LBS.

SULFENIMIDE FUNGICIDE

Active Ingredient	By Wt.
cis-N-[(1,1,2,2-tetrachloroethyl)thio]-4-cyclohexene-1,2-dicarboximide*	80%
Inert Ingredients	20%
*U. S. Pat. No. 3,178,447	

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE IN DRINKING WATER, FOOD OR DRINKING CONTAINERS.

BURN BOTTLES AND CONTAINERS. DO NOT REUSE.

STORE IN A COOL, DRY PLACE.
PROTECT FROM EXCESSIVE HEAT.

POTATOES: Early Blight, Late Blight - 2.0-2.5 quarts per acre in water to cover. Begin treatment when plants are 6" high and continue at 7-10 day intervals through severe disease conditions. A higher listed dosage is starter spray for 1st



READ THE LABEL

KEEP OUT OF REACH OF CHILDREN.

CAUTION: This product is highly toxic to fish and aquatic life. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption. Do not apply to plants or animals that are contaminated with insecticides, herbicides, or other pesticides.

COMPATIBILITY: This product is compatible with most commonly used insecticides, fungicides, and herbicides with the exception of those listed in the "Incompatibilities" section of the label. Do not apply to plants or animals that are contaminated with insecticides, herbicides, or other pesticides.

CONDITIONS OF SALE: This product is sold as a concentrate. It is not to be used as a spray. It is to be diluted with water before use. It is not to be used on plants or animals that are contaminated with insecticides, herbicides, or other pesticides. It is not to be used on plants or animals that are contaminated with insecticides, herbicides, or other pesticides. It is not to be used on plants or animals that are contaminated with insecticides, herbicides, or other pesticides.

CHEVRON CHEMICAL COMPANY

ORTHO DIVISION

San Francisco, California 94120

- RICHMOND, CALIFORNIA
- PORTLAND, OREGON
- FRESNO, CALIFORNIA
- DES MOINES, IOWA
- CHERRY HILL, NEW JERSEY
- DALLAS, TEXAS
- WHITTIER, CALIFORNIA
- HONOLULU, HAWAII
-
- ORLANDO, FLORIDA



237-1716



PEACHES, APRICOTS: Scab, Brown Rot, Brown Rot Blossom Blight, Whipscour Rot - 1 to 1 1/4 lbs. per 100 gals., using sufficient spray per acre, depending on tree size, to provide thorough coverage. Apply at prebloom, full bloom, petal fall and shoot split and repeat at 10 to 14-day intervals to the upper harvest. Do not apply after shoot split in Calif., Oregon, Washington or if fruit is to be dried.

PEACHES, NECTARINES, APRICOTS (Do not spray): Conium Blight (Shot Hole), Peach Leaf Curl - 2 lbs. per 100 gals. (or 8 lbs. per acre in water to cover), and apply in fall before rains begin and repeat in late winter prior to bud swell; or 4 lbs. per 100 gal. (or 16 lbs. per acre in water to cover) and apply as a single fall application prior to fall rains.

CHERRIES (sour): Cherry Leaf Spot - 1/2 to 1 lb. per 100 gals. spray, using sufficient spray per acre, depending on tree size, to provide thorough coverage. Begin sprays at petal fall and repeat at approximately 10-day intervals for 3 to 4 applications, making the last application to the foliage after harvest.

CHERRIES, PLUMS, PRUNES: Brown Rot Blossom Blight (California and Oregon Only) - 1 to 2 lbs. per 100 gals. using sufficient spray per acre, depending on tree size, to provide thorough coverage. Apply at prebloom, bloom and petal fall.

TOMATOES: Anthracnose, Early Blight, Late Blight, Gray Leaf Spot (Stemphylium), Hail Head Spot, Fruit Rot - 1 1/2 to 3 lbs. per acre in 100 to 150 gals. water. Apply when fruit of first clusters are well formed and repeat at 7 to 10-day intervals. When weather conditions favor disease development, use the higher rate and begin applications when plants are set in fields.

MELONS (Cantaloupe, Honeydew, Muskmelon & Others): Alternaria Leaf Blight, Downy Mildew, Anthracnose, Gummy Stem Blight - 1 1/2 to 3 lbs. per acre in 100 to 150 gals. spray as required for thorough coverage, depending on size of plants. Apply when first true leaves appear and repeat at weekly intervals. If powdery mildew is a problem, use a mildew fungicide as needed.

WATERMELONS: Gummy Stem Blight, Anthracnose, Downy Mildew - 1 1/2 to 3 lbs. per acre in 50 to 150 gals. spray as required for thorough coverage, depending on size of plants. Begin applications when plants start to vine and repeat at 5 to 10-day intervals as needed.

CUCUMBERS: Downy Mildew, Scab, Gummy Stem Blight, Anthracnose, Alternaria Leaf Blight - 1 1/2 to 2 lbs. per acre in 50 to 150 gals. of spray as required for thorough coverage, depending on size of plants. Apply when first true leaves appear and continue at weekly intervals. If Powdery Mildew is a problem, use a mildew fungicide as needed.