

G 3 Dust



ORTHO

239-1726

ACCEPTED

NOV 29 1972
UNITED STATES FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 239-1726

(INSECTICIDE)

Contains 3% GUTHION®

| | |
|---|--------|
| Active Ingredient | By Wt. |
| O,O-Dimethyl S-(4-oxo-1,2,3-benzotriazin-3(4H)-ylmethyl) Phosphorodithioate | 3% |
| Inert Ingredients | 97% |

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH DANGER STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER.

STORE IN COOL, DRY PLACE.

PROTECT FROM EXCESSIVE HEAT.

BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE.

DIRECTIONS

This product is formulated for application by ground or air equipment.

ALFALFA: Alfalfa Plant Bug, Aphids, Fleahoppers, Leafhoppers, Lygus, Mites, Spittlebug—20 to 25 lbs. per acre. Do not apply more than once between cuttings nor within 21 days of harvest. DO NOT USE IN CALIFORNIA.

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER: Aphids, Cabbage Looper, Imported Cabbage Worm, Diamond-back Moth—25 lbs. per acre. Repeat as necessary. Do not apply to Brussels Sprouts within 7 days of harvest; to Broccoli and Cauliflower within 15 days and to Cabbage within 21 days of harvest.

CLOVER: Aphids, Leafhoppers, Mites—15 to 25 lbs. per acre. Alfalfa Plant Bug, Fleahoppers, Lygus, Spittlebug—25 lbs. per acre. Do not apply more than once per cutting nor within 21 days of harvest.

NET WEIGHT 50 LBS.

Chevron Chemical Company

Ortho Division/San Francisco, Calif. 94119

Richmond, California Fresno, California Honolulu, Hawaii
Des Moines, Iowa Perth Amboy, New Jersey Orlando, Florida

Form 6505-1 Product 3284 Made in U.S.A.

EPA Reg. No. 239-1726-AA



POISON



ANTIDOTE: —If Swallowed. Give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Have victim lie down and keep quiet. **CALL A PHYSICIAN IMMEDIATELY!** If On Skin. In case of contact, remove contaminated clothing and immediately wash skin with soap and water.

DANGER:

KEEP OUT OF REACH OF CHILDREN

DO NOT TAKE INTERNALLY

DO NOT INHALE

DO NOT GET ON SKIN

DANGER: Poisonous If Swallowed, Inhaled or Absorbed Through Skin! Wear a mask or respirator of a type passed by the U. S. Bureau of Mines for Guthion protection. Do not get in eyes or on skin. Wear natural rubber gloves, protective clothing and goggles. In case of contact, wash immediately with soap and water. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use. Keep all unprotected persons out of operating areas or vicinity where there may be danger of drift. Vacated areas should not be re-entered for at least two days. **Note to Physician: Atropine is antidotal.** 2-PAM is also antidotal and may be administered in conjunction with atropine. Do not store or transport near feed or food. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

This product is toxic to fish, birds and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

GRAPES: Thrips, Grape ~~Leafhoppers, Mites,~~ Red-banded Leaf Roller—20 to 30 lbs. per acre. Do not apply more than 3 times per season. Do not apply within 10 days of harvest.

POTATOES: Aphids, Colorado Potato Beetle, Leafhoppers, Leaf Miners, Flea Beetle, Spittle Bug, Tarnished Plant Bug, Tuber Moth (West Coast Only)—25 lbs. per acre. Repeat as necessary. Do not apply within 7 days of harvest.

STRAWBERRIES: Meadow Spittle Bug, Oblique-banded Leaf Roller, Strawberry Leaf Roller—16 lbs. per acre. Do not apply within 5 days of harvest.

WALNUTS: Aphids, Codling Moth, European Red Mite—30 lbs. per acre. Repeat as necessary. Do not apply after husks split. Do not graze livestock in treated groves within 21 days after treatment.

CONDITIONS OF SALE: 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Chevron under 1 above) resulting from handling, storage, and use of this material.

GUTHION® trademark of Farbenfabriken Bayer, A. G. Chemagro Corporation licensee.

†U.S. Pat. No. 2,758,115—This material used under license from Chemagro Corporation