

# SEVIN\* Sulfur 7.5-75 Dust



# ORTHO

MAY 5 1972

239-174

## (INSECTICIDE-FUNGICIDE)

Active Ingredients	By Wt.
Carbaryl (1-naphthyl N-methylcarbamate)	7.5%
Sulfur	75.0%
Inert Ingredients	17.5%
U. S. Pat. Nos. 2,903,478-3,009,855	

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER.

BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE.

### DIRECTIONS

This product is formulated for application by ground or air equipment.

**BEANS:** Mexican Bean Beetle—10 lbs. per acre. Leafhoppers—15 lbs. per acre. Lygus, Stink Bugs, Powdery Mildew—25 lbs. per acre. Apply thoroughly at first signs of disease or insect infestation. When necessary repeat every 7 to 10 days. May be used up to harvest.

**CORN:** Corn Earworm—20 to 25 lbs. per acre. Apply at 3-day intervals starting when silks first appear. Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee populations. Make 4 to 5 applications until silks dry or turn brown. May be applied up to harvest.

**GRAPES:** Grape Leaf Folder—1st Brood: 15 to 20 lbs. per acre just before first brood larvae emerge from the first rolls. 2nd and 3rd Broods: 20 to 25 lbs. per acre. Grape Leafhopper, Powdery Mildew—15 lbs. per acre. Apply thoroughly.

## NET WEIGHT 50 LBS.

**Chevron Chemical Company**

Ortho Division San Francisco, Calif. 94119  
 Richmond, California Fresno, California Honolulu, Hawaii  
 Des Moines, Iowa Perth Amboy, New Jersey Orlando, Florida  
 Form 6516-1 Product 3286 Made in U.S.A.

oroughly at first signs of disease or insect infestation. When necessary repeat every 7 to 10 days. Thorough coverage is essential. May be applied up to harvest. This application may be expected to control DDT-resistant leafhoppers.

**COTTON:** Early-Season—Thrips, Cotton Fleahopper—10 to 12 lbs. per acre as soon as insects appear. Mid and Late-Season — Boll Weevil, Bollworm, Pink Bollworm, Cotton Leafworm, Lygus Bugs, Cotton Leaf Perforator—20 to 25 lbs. per acre. Apply when insects appear and repeat at 5 to 7-day intervals. Aphid populations will usually be suppressed when this insecticide is used on a regular schedule of treatments.

**PEACHES:** Oriental Fruit Moth, Peach Twig Borer, Plum Curculio, Cat-Facing Insects, Periodical Cicada, Powdery Mildew—40 to 45 lbs. per acre. Apply thoroughly at first signs of disease or insect infestation. When necessary repeat every 7 to 10 day. Do not apply within 1 day of harvest.

**POTATOES:** Colorado Potato Beetle—10 lbs. per acre. Leafhopper and Flea Beetle—15 lbs. per acre. Lygus Bugs—25 lbs. per acre. Apply when insects appear. Apply dust thoroughly to upper and lower leaf surfaces. If needed, repeat applications at 7 to 10-day intervals.

**TOMATOES:** Tomato Fruitworm, Fall Armyworm, Flea Beetles, Russet Mite, Tomato Hornworm, European Corn Borer, Stink Bugs—25 to 40 lbs. per acre. Apply thoroughly at first signs of insect infestation. Repeat as necessary. May be applied up to the day of harvest. Because of the sulfur content of this dust, first obtain permission from your canner or processor before using it on tomatoes intended for sale to such canner or processor.

**CONDITIONS OF SALE:** 1. Chevron Chemical Company Chevron warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability except those assumed by Chevron under 1 (above) resulting from handling, storage, and use of this material.



## READ THE LABEL

KEEP OUT OF REACH OF CHILDREN

**CAUTION:** HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN. Avoid Breathing Dust. Do Not Take Internally. Skin Contact May Be Harmful. Avoid Contact. Wash Hands and Face Before Eating. Take Shower or Bath After Work. Wear Regular Long Sleeve Work Clothing. Change to Clean Clothing Daily. **Note for Physician:** Carbaryl is a moderate, reversible, Cholinesterase inhibitor. Atropine is antidotal. Do not store or transport near feed or food. AVOID CONTAMINATION OF FOOD, FEED, FEEDING TROUGHS AND WATERING RECEPTACLES. Some fruits and other plants are susceptible to injury from sulfur under certain climatic conditions. The user is advised not to use sulfur on any crop unless local use has proved that sulfur does not damage crops in that locality. **Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.**

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension