

[ 239-1782 ]



# Malathion - Cryolite 4-35 Dust

[INSECTICIDE]

NET WEIGHT 50 LBS.

Active Ingredients	By Wt.
Malathion	4%
Sodium Fluoaluminat	35%
Inert Ingredients	61%
O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate.	
Fluorine expressed as elemental	19%

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER.

BE CAREFUL WHEN APPLYING NEAR OPENINGS OF SMOKE.

This product is formulated for application by ground or air equipment.

Apply thoroughly. Do not apply on food crops within 3 days of harvest unless otherwise specified. Remove residues remaining at harvest by washing, brushing or other effective means.

**CUCUMBERS, SQUASH, MELONS:** Aphids, Spider Mites, Pickleworm, Cucumber Beetles, Flea Beetles; **CANTALOUPE** — Leafhoppers — 30 to 35 lbs. per acre. Repeat as necessary. **SQUASH:** Squash Vine Borers—40 lbs. per acre. **PUMPKINS:** Squash Vine Borers—45 lbs. per acre. Apply twice a week to stems and vines at base of plants. Do not apply to Pumpkins within 3 days of harvest, and to Cucumbers, Cantaloupes, Squash and Melons within 1 day of harvest. Do not apply unless plants are dry.



READ THE LABEL

KEEP OUT OF REACH OF CHILDREN

**CAUTION:** Avoid breathing dust. Avoid contact with skin; wash thoroughly after using. Do not store or transport near feed or food. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.


This product will kill fish. Keep out of any body of water. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

**LETTUCE:** Aphids, Mites, Cabbage Looper, Leafhoppers, Climbing Cutworms, Alfalfa Looper, Corn Earworm—30 to 40 lbs. per acre. Do not apply after the seedling stage.

**TOMATOES:** Aphids, Spider Mites, Tomato Russet Mite, Drosophila, Armyworms, Flea Beetles—35 to 45 lbs. per acre. Do not apply within 1 day of harvest.

**CONDITIONS OF SALE:** 1. Chevron Chemical Company Chevron warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risk and liability, except those assumed by Chevron under 1 above, resulting from handling, storage and use of this material.



## CHEVRON CHEMICAL COMPANY

FORM 1603 E

USDA Reg. No. 239-1782

ORTHO DIVISION — San Francisco, California 94120

Product 3335

Made in U. S. A.

RICHMOND, CALIFORNIA

PORTLAND, OREGON

FRESNO, CALIFORNIA



DES MOINES, IOWA

CHERRY HILL, NEW JERSEY

HONOLULU, HAWAII

ORLANDO, FLORIDA