

Malathion 6 Dust



ORTHO

[239-1796]

(INSECTICIDE)

Active Ingredient Malathion* By Wt. 6%
Inert Ingredients 94%
*O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER.

EMPTY BAGS IMMEDIATELY WHEN EMPTY. STAY OUT OF BUCKETS.

Apply to plants, soil, or other surfaces where insects or mites are present. Do not apply to plants or soil that are being prepared for planting. Do not apply to plants or soil that are being prepared for planting.

ALFALFA: Armyworms, Aphids, Potato Leafhoppers, Spider Mites, Alfalfa Weevil Larvae, Spittlebugs, Vetch Bruchids—20 lbs. per acre. Spotted Alfalfa Aphids—Ground—10 to 12 lbs. per acre. Air—10 to 16 lbs. per acre. Apply to alfalfa before cutting, or to the cutting or early in the morning when the dew is still on the leaves. Do not apply to alfalfa that is being cut for hay. May be applied on day of harvest or grazing.

APPLES: Red Spider Mites (such as European Red Mite, Two-Spotted Mite, Pacific Mite, Willamette Mite and Clover Mite), Green Apple Aphid, Rosy Apple Aphid, Woolly Apple Aphids—20 to 40 lbs. per acre. This product may be applied to apples during the growing season.

APRICOTS: Spider Mites, Brown Apricot Scale (Crawlers), Orange Tortrix—50 lbs. per acre. Do not apply within 7 days of harvest.

ASPARAGUS: Thrips—20 lbs. per acre. Repeat as necessary. Do not apply within 7 days of harvest.

BEANS: Mexican Bean Beetle, Japanese Beetles, Cucumber Beetles, Leafhopper, Spider Mites, Aphids—30 lbs. per acre. Two or more applications as needed. Do not apply within 1 day of harvest.

BLUEBERRIES: Japanese Beetles—20 lbs. per acre. Apply when berries turn blue and are 10 days before harvest. Do not apply to Plum Curculio, Sharp-nosed Leafhopper—30 to 40 lbs. per acre. Do not apply within 1 day of harvest.

BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, CABBAGE, KALE, MUSTARD GREENS, TURNIPS: Aphids, Imported Cabbageworm, Flea Beetles—30 to 40 lbs. per acre. Repeat as necessary. Do not apply to Cauliflower, Broccoli, Sprouts, Cabbage, Mustard Greens and Kale within 7 days of harvest and to Broccoli and Turnips within 3 days of harvest.

NET WEIGHT 50 LBS.

Chevron Chemical Company

Ortho Division/San Francisco, Calif. 94119

Richmond, California Fresno, California Honolulu, Hawaii
Des Moines, Iowa Perth Amboy, New Jersey Orlando, Florida

Form 6658-G EPA Reg. No. 239 1796 Product 3361 Made in U.S.A.

CARROTS: Leafhoppers—30 lbs. per acre. Repeat as necessary. Do not apply within 7 days of harvest.

CLOVER: Aphids, Mites—18 to 20 lbs. per acre. Do not apply to Clover in bloom. May be applied on day of harvest or grazing.

COTTON: Aphids, Spider Mites—25 to 30 lbs. per acre. Two or more applications as needed.

CRANBERRIES: Leafhoppers, Black-Headed Fireworms—Ground Machine: 20 lbs. per acre. Make thorough coverage.

CUCUMBERS, SQUASH, MELONS: Aphids, Spider Mites, Pickleworms, Squash Vine Borer—20 to 35 lbs. per acre. **CANTALOUPE—**Leafhoppers—30 to 35 lbs. per acre. Repeat as necessary. Do not apply unless plants are dry and without heavy dew.

DATES: Nitidulid Beetles—Apply to trees on each side of thoroughly watered 20 to 30 lb. per acre. Make no application on dates within 7 days of the first picking.

FILBERTS: Filbert Aphid, Apple Mealybug, Eye-spotted Bud Moth—40 lbs. per acre. Repeat as necessary.

GRAPES: Leafhoppers, Spider Mites—15 to 30 lbs. per acre. Leafhoppers—One or more applications as needed. Spider Mites—Two or more applications as needed.

LETTUCE: Aphids, Mites, Cabbage Looper, Leafhoppers—25 to 30 lbs. per acre. Do not apply to Head Lettuce within 7 days of harvest and to Leaf Lettuce within 14 days.

ONIONS: Onion Thrips, Onion Maggots—30 lbs. per acre. Do not apply to onions within 3 days of harvest.

ORNAMENTALS: Aphids, Red Spider Mites, Whitefly, Thrips, Japanese Beetle Adult, Four-lined Leaf Bug, Bagworm, Tarnished Plant Bug, Rose Leafhopper, Potato Leafhopper—25 to 30 lbs. per acre. Repeat as necessary. Do not apply to Begonia, Madia and Pteris ferns, and some species of Crataegus.

PEARS: Pear Psylla, Aphids, Woolly Aphids—30 to 40 lbs. per acre. One or more applications as needed. Spider Mites—Same directions as for Apples. Do not apply within 1 day of harvest.

ACCEPTED
May 11 1972
UNDER THE FEDERAL INSECTICIDE ACT
FOR THE UNITED STATES DEPARTMENT OF AGRICULTURE
WASHINGTON, D.C.

PEAS: Pea Aphid—20 lbs. per acre as needed. Do not apply within 3 days of harvest or 7 days if forage is to be fed to livestock.

PEPPERMINT, SPEARMINT: Flea Beetle Adults—17 lbs. per acre. Do not apply within 7 days of cutting.

PEPPERS: Pepper Maggots—24 lbs. per acre.

POTATOES: Aphids, Leafhoppers—25 to 30 lbs. per acre. One or more applications as needed. Use up to harvest.

PUMPKINS: Squash Vine Borers—33 lbs. per acre. Apply twice a week to main and vine at base of plants. Do not apply unless plants are dry.

SPINACH: Aphids—25 to 30 lbs. per acre. Do not apply within 7 days of harvest.

RASPBERRIES: Aphids, Rose Scale—35 lbs. per acre. Repeat as necessary. Do not apply within 1 day of harvest.

STRAWBERRIES: Aphids, Spider Mites, Lygus Bugs, Spittlebugs, Potato Leafhoppers, Strawberry Leafrollers, Whiteflies, Strawberry Root Weevil, Field Crickets—20 lbs. per acre. Repeat as necessary. Strawberry Root Weevils—Apply to plants during planting and work into top 6 to 8 inches of soil.

TOBACCO: Aphids—20 to 30 lbs. per acre. Repeat as necessary. Do not apply within 7 days of harvest.

TOMATOES: Aphids, Spider Mites, Tomato Russet Mite, Drosophila—20 to 40 lbs. per acre. Do not apply within 5 days of harvest.

MILKWEEDS: ... Around C ... Around

DOGWOOD: ... Around C ... Around

DOGWOOD: ... Around C ... Around

DOGWOOD: ... Around C ... Around

DOGWOOD: ... Around C ... Around

DOGWOOD: ... Around C ... Around

DOGWOOD: ... Around C ... Around

DOGWOOD: ... Around C ... Around

DOGWOOD: ... Around C ... Around

DOGWOOD: ... Around C ... Around

DOGWOOD: ... Around C ... Around

DOGWOOD: ... Around C ... Around

DOGWOOD: ... Around C ... Around

DOGWOOD: ... Around C ... Around

DOGWOOD: ... Around C ... Around

DOGWOOD: ... Around C ... Around

DOGWOOD: ... Around C ... Around

CAUTION

READ THE LABEL

KEEP OUT OF REACH OF CHILDREN

CAUTION: Avoid breathing dust. Avoid contact with skin. Wash thoroughly after using. Do not store or transport near food or feed. Do not apply under conditions involving possible drift to food, storage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

Highly toxic to fish. Do not contaminate any body of water nor apply where conditions favor drift from the crop area treated. This product is highly toxic to birds exposed to direct treatment or residues on crops. Protective instructions may be obtained from your Cooperative Agricultural Extension Office.