

eyes Avoid wash eyes with on skin wash person to Call a doctor mation call m liquid and an this product is actively visit away from se this product WITH PRECAU TM APPLICABLE early storage water that be disposed of then dispose of iprocedures

Berry Moth, Grape Leaf Skeletonizer, Japanese Beetle, Leafhoppers, Rose Choler, San Jose Scale (Crawlers), BLACKBERRIES, RASPBERRIES Japanese Beetle, Leafhoppers, Rose Choler, San Jose Scale (Crawlers) **STRAWBERRIES** Japanese Beetle, Leafhoppers, Rose Choler, Spotted Wing, Strawberry Weevil. 2 to 3 qts. per 100 gas water. Apply at regular intervals and repeat at 7 to 14 day intervals as needed. Do not apply to Blueberries, Cranberries, Currants, Gooseberries and Grapes within 14 days of harvest. Do not apply to Raspberries, Logberries, Raspberries and Strawberries within 14 days of harvest. However, application may be made up to 3 days of harvest at a rate of 1/2 lbs. actual Methoxychlor per acre.

GARDEN CROPS: ASPARAGUS - Asparagus Beetle, Fall Armyworm, Flea Beetle, Japanese Beetle, BEANS, BLACKEYED PEAS, PEAS - Bean Leaf Beetle, Fall Armyworm, Flea Beetle, Japanese Beetle, Leafhoppers, Mexican Bean Beetle, Pea Weevil, BEETS - Fall Armyworm, Japanese Beetle, BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULI FLOWER, RALE, LETTUCE, BUTABAGAS, SPINACH, COLLARDS, KOLH, RAM TURNSIPS - Imported Cabbageworm, Fall Armyworm, Flea Beetle, Japanese Beetle, CANNONTS, RADISHES - Fall Armyworm, Flea Beetle, Japanese Beetle, CORN - Fall Armyworm, Flea Beetle, Japanese Beetle, Leafhoppers, CANTALOUPE, CUCUMBERS, MELONS, PUMPKINS, SQUASH, SUMMER SQUASH - Blosser Beetle, Cucumber Beetle, Fall Armyworm, Flea Beetle, Japanese Beetle, Leafhoppers, Squash Vine Borer, EGGPLANT, TOMATOES, PEPPERS - Blosser Beetle, Colorado Potato Beetle, Fall Armyworm, Flea Beetle, Japanese Beetle, Leafhoppers, POTATOES - Colorado Potato Beetle, Flea Beetle, Japanese Beetle, Leafhoppers. 2 to 3 qts. per 100 gas water. Apply at regular intervals and repeat at 7 to 14 day intervals as needed. Do not apply to corn, melons, peas, radishes and rutabagas within 14 days of harvest. Do not apply to beets, broccolis, brussels sprouts, cauliflowers, collards, kohlrabi, lettuce, spinach and turnsips within 14 days of harvest. Asparagus beetle may be controlled by washing plants with water at harvest. Must be applied before harvest. Do not apply to beets, eggplant, radishes, peppers, pumpkin, squash and tomatoes up to 14 days of harvest at a rate of 1/2 lbs. actual Methoxychlor per acre. Maximum of 2 applications per season.

ELMS - Elm Bark Beetle Control - To prevent transmission of Dutch Elm Disease. Hydraulic Sprayer Application - 2 qts. per 100 gas water. Spray from July to October. Mist Blower Application - 1 qt. per 100 gas water. Apply from July to September. Do not apply to elms that are above 40 ft. or in the shade.

ALFALFA, CLOVER, GRASS, SOYBEANS, COMPEAS, PEANUTS, FORAGE CROPS - Leafhoppers, Spotted Wing, Alfalfa Caterpillar, Alfalfa Weevil, Leafhoppers, Alfalfa Weevil, Loras, Alfalfa Weevil, Fall Armyworm, Clover Leaf Weevil, Mexican Bean Beetle, Pea Weevil, Velvet Bean Caterpillar, Alfalfa Caterpillar, Compeas Caterpillar, Japanese Beetle, Leafhoppers, Fall Armyworm. 2 to 3 qts. per 100 gas water. Apply at regular intervals and repeat at 7 to 14 day intervals as needed. Do not apply to alfalfa, clover, grass, soybeans, compeas, peanuts, forage crops within 14 days of harvest.

GRAIN STORAGE BINS - To aid in Control of Corn Ear Moth, Lesser Grain Borer, Lesser Grain Weevil, Red Flour Beetle, Long Headed Flour Beetle, Rice Weevil, Grainy Weevil, Confused Flour Beetle, Foreign Grain Beetle, Hairy Fungus Beetle. 1 qt. per 100 gas water. Apply at regular intervals and repeat at 7 to 14 day intervals as needed. Do not apply to bins containing grain within 14 days of harvest. NOTE: Do not apply to bins containing grain within 14 days of harvest.

CONTACT AND SPACE SPRAY FOR HOUSE FLIES AND STABLE FLIES in Farm Buildings, Barnyards, Alleys, on Dumps and Other Outdoor Locations. Dilute 1 qt. in 2 1/2 gal. water and apply with a backpack sprayer or directly to flies by hydraulic sprayer. Do not apply to human or animal handling equipment. Care should be taken to avoid contamination of the water supply. Flies should not be present when animals are present in the vicinity.

LIVESTOCK (except Dairy Animals and Poultry) - Use as spray only. Do not use in dipping vats.

CATTLE 1 qt. in 12 1/2 gal. water. Apply as a spray to the head, neck and control of Horn Flies. For Control of Short Nosed and Long Nosed Cattle Lice. Apply as a drenching spray. Hogs, Lice. 2 to 3 qts. per 100 gas water. Apply as a drenching spray.

SELF-APPLICATOR FLY CONTROL ON CATTLE (except Dairy Animals) Horn Flies. 1 qt. in 12 1/2 gal. water. Apply as a spray to the head, neck and control of Horn Flies. For Control of Short Nosed and Long Nosed Cattle Lice. Apply as a drenching spray. Hogs, Lice. 2 to 3 qts. per 100 gas water. Apply as a drenching spray.

Do not apply under conditions involving possible dirt to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

COMPATIBILITY - Do not mix with other agricultural chemicals. Do not mix with other insecticides.

IF EXPOSED TO PROLONGED COLD, PLACE IN WARM STORAGE (50° to 60° F) FOR SEVERAL HOURS AND AGITATE BEFORE USING

CONDITIONS OF SALE - This material is sold as a material only. It is not to be used in any other manner. The material is sold as a material only. It is not to be used in any other manner. The material is sold as a material only. It is not to be used in any other manner.



ORTHO

Methoxychlor 2 Emulsive

(INSECTICIDE)

ACCUMULATED
APR 13 1982

**KEEP OUT OF REACH OF CHILDREN
WARNING
FOR AGRICULTURAL USE ONLY DO NOT STORE IN
AREAS ACCESSIBLE TO CHILDREN.
SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY
STATEMENTS.**

Chevron Chemical Company
1800 Avenue of the Stars
P.O. Box 988
Houston, Texas 77268
U.S.A.

NET CONTENTS 1 QUART