

# ORTHO<sup>®</sup>

[INSECTICIDE]

NET CONTENTS  
LIQUID

**XYLENE BASE SOLVENT**

Active Ingredients	By Wt.
Toxaphene*	58%
Xylene Base Aromatic Petroleum Solvent	38%
Inert Ingredients	4%
*Technical Chlorinated Camphene (Chlorine Content 67% to 69%) Contains 6 lbs. Toxaphene per gallon.	

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

WASH AND DESTROY CONTAINER WHEN EMPTY. NEVER RE-USE.


### DIRECTIONS

USE DESIGNATED AMOUNT OF PRODUCT IN THE FOLLOWING GALLONS WATER PER ACRE UNLESS OTHERWISE NOTED:

GROUND APPLICATION—Vegetable and Field Crops—30 to 250 gals. Cotton—3 to 20 gals.  
Where Recommended for AIR APPLICATION—Field Crops—3 to 10 gals.

**BROCCOLI, CAULIFLOWER, CABBAGE, BRUSSELS SPROUTS:** Cabbage Caterpillars, Cabbage Looper—1 quart per acre in sufficient water to cover. Do not apply to Broccoli, Brussels Sprouts and Cauliflower after edible parts begin to form or within 30 days of harvest. Do not apply to Cabbage after heads start to form.

ACCEPTED  
100 23, 110  
239-18873  
UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND ROENTGENIC CHEMICALS ACT, REGISTERED TO ATTACHED COMMENTS.



**DO NOT INHALE**  
**DO NOT GET ON SKIN**  
**DO NOT TAKE INTERNALLY**  
**KEEP OUT OF REACH OF CHILDREN**

### WARNING:

May Be Fatal if Swallowed! Do not breathe spray mist. Do not get in eyes, on skin or on clothing. Wash thoroughly after using. Keep away from children and domestic animals. Do not store or transport with feed or food. **DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.** Do not use on young plants. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

Highly toxic to Fish and Wildlife: Birds feeding on treated areas may be killed. Do not contaminate any body of water or apply to areas or crops other than those for which directions are given on the label. This product is toxic to bees and should not be applied when bees are actively visiting the area.

**CORN:** Armyworms, Grasshoppers, Chinch Bugs—Apply 1½ qts. per acre in sufficient water for coverage. Apply as first insects appear. Repeat as necessary. Do not feed treated forage to dairy animals or animals being finished for slaughter.

**COTTON:** Cotton Boll Weevil, Cotton Bollworm, Armyworm, Fleahopper, Thrips, Leafworm, Grasshoppers—Use 1½ quarts per acre diluted in sufficient water to give even coverage. By Airplane—3 to 5 gallons water per acre. By Ground Rig—5 to 15 gallons water per acre. For Early Application for Control of Thrips and Fleahoppers only: Use ¾ quart diluted in sufficient water to insure even coverage, depending on method of application. Do not graze dairy animals or animals being finished for slaughter in fields treated late in the season.

**LETTUCE:** Cutworms (Surface Feeding and Climbing)—1 to 1½ qts. per acre in water to cover. Do not apply after seedling stage on Leaf Lettuce. Do not apply to head lettuce after heads begin to form.

**PEPPERS:** Cutworms—2 qts. per acre in water to cover. Do not apply within 5 days of harvest.

**SOYBEANS:** Velvetbean Caterpillar, Bollworm, Corn Earworm, Bean Leaf Beetle, Armyworms, Grasshoppers, Blister Beetle, Green Cloverworm—1½ to 2½ quarts per acre in water to cover. Repeat as necessary. Do not apply less than 3 weeks before harvest. Do not feed plants treated with Toxaphene or ensilage made from treated plants to poultry, dairy animals or animals being finished for slaughter. Do not feed soybean mill trash to livestock or poultry.

### LIVESTOCK

Do not treat Beef Cattle, Goats, Sheep or Swine within 28 days of slaughter. Do not use on dairy animals or in dairy barns. Do not treat calves under 3 months of age.

**SHEEP TICKS:** Sheep—As a Dip—1½ quarts in 100 gals. water. As a Spray—3 quarts in 100 gals. water, spraying thoroughly during warm weather, when wool is only a few inches long.

**LICE:** Beef Cattle, Swine—As a Spray—3 quarts in 100 gals. water. Do not use on Dairy Cattle. Sheep, Goats—As a Dip—1 quart in 100 gals. water. Do not use on Dairy Goats.

**HORN FLIES:** Beef Cattle—As a Spray—3 quarts in 100 gals. water, applying about 2 quarts per mature animal. Do not use on Dairy Cattle.

**TICKS:** Beef Cattle, Swine—As a Spray—3 quarts in 100 gals. water. Beef Cattle—As a Dip—1½ quarts in 100 gals. water. Do not dip heavy, fat cattle in hot weather. Do not use on Dairy Cattle.

**CABLE TYPE BACK RUBBERS FOR Range Cattle to Control Hornflies, Mosquitoes**—¾ pint ORTHO Toxaphene 6 Emulsive XB to 1 gallon No. 2 fuel oil or furnace oil. Do not use on dairy animals. Do not treat Beef Cattle within 28 days of slaughter.

**CONDITIONS OF SALE:** 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Chevron under 1. above) resulting from handling, storage, and use of this material.