

[INSECTICIDE]

NET WEIGHT 50 LBS.

Active Ingredients By Wt
 Carbophenothion*** 2%
 Carbaryl 1-naphthyl N-methylcarbamate 5%
 Inert Ingredients 93%
 ***S-(p-chlorophenylthio) methyl O,O-diethyl phosphorodithioate
 U. S. Pat. Nos. 2,793,224-2,903,478-3,000,855

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS

KEEP OFF OF VEGETATION AND OTHER SURFACES TO BE TREATED UNTIL DRY. DO NOT ALLOW CHILDREN OR OTHERS TO TOUCH TREATED SURFACES.

DIRECTIONS

APPLES, PEARS: Apple Aphid, Coding Moth, European Red Mite, Rosy Apple Aphid, Two-Spotted Mites, Woolly Apple Aphid, Green Apple Aphid, Red-banded Leaf Roller, Pear Psylla, Periodical Cicada, Apple Maggot.

CHERRIES, PEACHES, PLUMS, PRUNES: Aphids, Orchard Mite, Oriental Fruit Moth, Peach Twig Borer, Periodical Cicada, Coding Moth, Eastern Tent Caterpillar, Orange Tortrix, Western Tussock Moth, Cherry Fruit Fly.

Apply to foliage and fruit. Do not apply to fruit directly. Repeat applications as needed. Do not apply with rain or irrigation.




READ THE LABEL

KEEP OUT OF REACH OF CHILDREN

CAUTION: HAZARDOUS TO HEALTH. Avoid breathing dust. Wear a respirator and goggles. Avoid contact with skin and eyes. Wash hands and face thoroughly with soap and water after use. Do not eat, drink, or smoke while using. Do not apply under conditions involving possible drift to food, forage or other plants that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

CONDITIONS OF SALE

Product is sold as a concentrate. It must be diluted before use. Do not use in areas where it may be harmful to humans, animals, or the environment.



CHEVRON CHEMICAL COMPANY

ORTHO DIVISION — San Francisco, California 94120

FORM 6343 H

USDA Reg. No. 239-1997

Product No. 3107

Made in U.S.A.

RICHMOND, CALIFORNIA

PORTLAND, OREGON

FRESNO, CALIFORNIA

DES MOINES, IOWA

CHERRY HILL, NEW JERSEY

DALLAS, TEXAS

HONOLULU, HAWAII

ORLANDO, FLORIDA



®

ORTHO TRITHION* SEVIN** 2-5 Dust

[INSECTICIDE]

NET WEIGHT 50 LBS.

Active Ingredients By Wt
 Carbophenothion*** 2%
 Carbaryl (1-naphthyl N-methylcarbamate) 5%
 Inert Ingredients 93%
 ***S-[p-chlorophenylthio] methyl O,O-diethyl phosphorodithioate
 U. S. Pat. Nos. 2,793,224-2 503,478-3,009,855

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINERS

BUY IN BAG IMMEDIATELY AFTER EMPTYING CONTAINER

DIRECTIONS

Apply when insects are present. Keep in dust form. Formulated for use in dust form. Thorough coverage of plants with insecticide is essential.

APPLES, PEARS: Apple Aphid, Codling Moth, European Red Mite, Rosy Apple Aphid, Two-Spotted Mites, Woolly Apple Aphid, Green Apple Aphid, Red-banded Leaf Miner, Pear Psylla, Periodical Cicada, Apple Maggot. Apply 10-25 lbs. per acre. Repeat at 10 day intervals. Do not apply within 30 days of harvest.

CHERRIES, PEACHES, PLUMS, PRUNES: Aphids, Orchard Mite, Oriental Fruit Moth, Peach Twig Borer, Periodical Cicada, Codling Moth, Eastern Tent Caterpillar, Orange Tortrix, Western Tussock Moth, Cherry Fruit Fly. Apply 10-25 lbs. per acre. Repeat at 10 day intervals. Do not apply within 30 days of harvest.



READ THE LABEL

KEEP OUT OF REACH OF CHILDREN

CAUTION: Avoid contact with skin, eyes, nose, mouth, and clothing. Avoid contact with food, feed, or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.


GRAPES: Grape Leaf Folder, Grape Leafhopper, Two-Spotted Mites. Thorough coverage is essential. This material is not effective against the common DDT resistant leafhoppers which cause damage 30 days of harvest.

SNAP BEAN, LIMA BEAN, SOYBEAN (in pods): Bean Aphids, Mexican Bean Beetle, Serpentine Leaf Miner, Potato Leafhopper, Two-Spotted Mites, Lygus and Stink Bugs. 40 lbs. per acre. Do not apply within 7 days of harvest. Do not graze or feed crop material to livestock.

BEST DOCUMENT AVAILABLE

CONDITIONS OF SALE: Chevron Chemical Company, Chevron, warrants that this material conforms to the chemical description on the label and is reasonably fit for use in directions thereon. It does not make any guarantee, warranty, or representation, express or implied, as to the fitness or of MERCHANTABILITY of this material for any purpose other than that intended by the manufacturer.

It is the policy of Chevron Chemical Company to provide the best possible product. However, Chevron's control over the material is limited to the extent of its contribution with the manufacturer. Such risks include, but are not limited to, the possibility of drift to other plantings caused by drift to other plantings. Such risks occur even though the product is applied in accordance with the uses stated hereon and even though the manufacturer, Chevron, and user acknowledge and assume responsibility for the material as furnished by Chevron under the conditions stated hereon. No liability shall be assumed by Chevron for any damage or injury caused by the use of this material.



CHEVRON CHEMICAL COMPANY

FORM 6343 H USDA Reg. No. 239-1997 ORTHO DIVISION - San Francisco, California 94120 Product No. 3207 Made in U. S. A.

RICHMOND, CALIFORNIA PORTLAND, OREGON FRESNO, CALIFORNIA DES MOINES, IOWA CHERRY HILL, NEW JERSEY DALLAS, TEXAS
 HONOLULU, HAWAII