

# ORTHO<sup>®</sup> Sulfur 15-65 Dust



# ORTHO

## (FUNGICIDE)

Active Ingredients	By Wt.
Captan*	15%
Sulfur	65%
Inert Ingredients	20%
*N-(trichloromethylthio)-4-cyclohexene-1,2-dicarboximide.	

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER.

STORE IN COOL, DRY PLACE  
PROTECT FROM EXCESSIVE HEAT

BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE.

### DIRECTIONS

This product is formulated for application by ground or air equipment.

**CELERY:** Pink Rot, Damp-Off—(Field)—13 lbs. per acre. (Seed Bed)—25 to 30 lbs. per acre. Apply when all seeds have germinated. Repeat as necessary.

## NET WEIGHT 50 LBS.

### Chevron Chemical Company

Ortho Division/San Francisco, Calif. 94119  
 Richmond, California    Fresno, California  
 Des Moines, Iowa    Perth Amboy, New Jersey    Orlando, Florida  
 Form 6. '8-G    Product 3598    Made in U.S.A.  
 EPA Reg. No. 239-2013-AA

**GRAPES:** Botrytis Rot (Bunch Rot), Powdery Mildew, Silver Mite, Storage Rots—33 lbs. per acre. Make 2 applications before bloom and 1 application just after bloom. Repeat periodically making up to 3 cover applications before bunches close.

**PEACHES:** Brown Rot, Storage Rots, Powdery Mildew, Silver Mite, Peach Scab—33 lbs. per acre. Make 2 to 3 prebloom applications. Repeat at petal fall and 3 to 4 covers as necessary. Make preharvest application where necessary.

**PLUMS:** Brown Rot, Storage Rots, Powdery Mildew, Silver Mite—33 lbs. per acre. Make 2 to 3 prebloom applications. Repeat at petal fall and 3 to 4 covers as necessary. Make preharvest application where necessary.

**POTATOES:** Early and Late Blight—Begin applications when

plants are 2 to 4 inches high and repeat at 7 to 10-day intervals as long as disease threatens. Apply 15 to 20 lbs per acre.

**ROSES (Field):** Rust, Powdery Mildew—35 to 40 lbs. per acre. Apply to give even coverage of upper and lower leaf surfaces at first sign of disease. Repeat at regular intervals as necessary to maintain control.

**STRAWBERRIES:** Botrytis Rot, Powdery Mildew—20 lbs. per acre immediately after picking. Repeat applications after picking as necessary.

**TOMATOES (Established Plants):** Early Blight, Late Blight, Anthracnose, Russet Mite—On young plants, apply 15 to 20 lbs. per acre, increasing to 25 lbs. per acre on mature plants. Apply at first sign of insects and diseases. Repeat at 5 to 7-day intervals as necessary for control. Because of the sulfur content of this dust, first obtain permission from your canner or processor before using it on tomatoes intended for sale to such canner or processor.

**COMPATIBILITY:** Do not use this product on foliage closely following oil sprays.

**CONDITIONS OF SALE:** 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty, of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material.

2. Critical and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Chevron under 1 above) resulting from handling, storage, and use of this material.



## READ THE LABEL

### KEEP OUT OF REACH OF CHILDREN

**CAUTION:** Avoid inhalation of dust. Avoid contact with skin. Do not store or transport near feed or food. Some fruits and other plants are susceptible to injury from sulfur under certain climatic conditions. The user is advised not to use sulfur on any crop unless local use has proved that sulfur does not damage crops in that locality. **Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.**

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from areas treated.

ACC