

Endrin 1.6 Emulsive

ACCEPTED
MAY 16 1975
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 215-213



ORTHO

'PAEs' 239-MO-1 No. 000743 AA

(INSECTICIDE-RODENTICIDE)

| Active Ingredients | By Wt. |
|---|--------|
| Endrin | 18.6% |
| Petroleum hydrocarbons | 73.0% |
| Inert Ingredients | 8.4% |
| *Hexachloroepoxyoctahydro-endo,endo-dimethanonaphthalene. Contains 1.6 lbs. Endrin per gallon. | |

DANGER: KEEP OUT OF REACH OF CHILDREN.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH DANGER STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP OFFICE FLOOR CLEAN. CONTAINER IS NOT TO BE RE-USE. ... **DO NOT USE FOR OTHER PURPOSES.**

ORTHO Endrin 1.6 Emulsive

USE DESIGNATED AMOUNT OF PRODUCT IN THE FOLLOWING GALLONS WATER PER ACRE UNLESS OTHERWISE NOTED.

| | |
|------------------------------------|----|
| GROUND APPLICATION - Field Crops | 10 |
| Hand Gun Application - Cotton | 5 |
| Hand Gun Application - Field Crops | 10 |

When Recommended for ATR APPLICATION - Field Crops:

NET CONTENTS 5 GALS. LIQUID

Chevron Chemical Company
San Francisco, CA 94114 Richmond, CA 94804
© 1975

COTTON: Thrips, Cotton Fleahoppers, Rapid Plant Bug, Tarnished Plant Bug—Early and Mid-Season Control: 1 gal. in water to cover 10 acres. Late Season Control: 1 gal. in water to cover 8 acres. Early Season Boll Weevil and Bollworm (not Pink Bollworm) Control: Use 1 gal. in water to cover 8 acres. Mid-Season and Late Season Control of Bollworm and Boll Weevil: Use 1 gal. in water to cover 4 to 8 acres in compliance with other recommendations. Grasshoppers: 1 gal. in water to cover 8 acres.

DANGER:
DO NOT INHALE
DO NOT GET ON SKIN
DO NOT TAKE INTERNALLY

POISON

ANTIDOTE: IF SWALLOWED: Give a Tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Have victim lie down and keep quiet. **CALL A PHYSICIAN IMMEDIATELY.** Note to Physicians: Emergency Information - call 415-288-7500. There is no specific antidote for Endrin. Treat symptomatically. In case of poisoning administer symptomatic treatment as outlined above.

IF ON SKIN: Wash immediately with soap and water.

DANGER: ... **POISON.** ... **ANTIDOTE:** ... **IF ON SKIN:** ... **IF SWALLOWED:** ... **CONDITIONS OF SALE:** ... **NOTE:** ... **USE:** ... **REMARKS:** ...

Cabbage Looper, Celery Leaf Tier, Leaf Perforator, Salt Marsh Caterpillar, Brown Cotton Leafworm, Cotton Leafworm, Lygus Bugs, Cutworms — 1 gal. per acre in water for coverage. Do not apply within 5 days of fruit picking. Workers entering the field within 5 days after treatment should be protected. Do not get on bare skin. ... **ORTHONALS:** Commercial Plantings such as Cyclamen, Azaleas, Fuchsia, Gloxinia and English Ivy: Cyclamen Mite — 1 gal. per acre in water. ... **MOUSE CONTROL IN APPLE ORCHARDS:** Postharvest application to orchard floor in Oct. or Nov. Pine Mice — 1 gal. per acre in water. Boom Application: ... Hand Gun Application: ...

ORTHONALS: Commercial Plantings such as Cyclamen, Azaleas, Fuchsia, Gloxinia and English Ivy: Cyclamen Mite — 1 gal. per acre in water. ... **MOUSE CONTROL IN APPLE ORCHARDS:** Postharvest application to orchard floor in Oct. or Nov. Pine Mice — 1 gal. per acre in water. Boom Application: ... Hand Gun Application: ...

MIXING: ... **CONDITIONS OF SALE:** ... **NOTE:** ... **USE:** ...

CONDITIONS OF SALE: ... **NOTE:** ... **USE:** ...

NOTE: ... **USE:** ... **REMARKS:** ...