

ORTHO

NET WEIGHT 50 LBS.

[INSECTICIDE]

Active Ingredient	By Wt.
O,O-dimethyl O-p-nitrophenyl thiophosphate	25%
Inert Ingredients	75%

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS, WARNINGS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

Containers should be disposed of by burying with wastes or by burning. (Keep out of smoke.)

Not for use or storage in or around the home.

NOTE—

For dilute (ground) application on all crops: Use recommended amounts in 30-100 gallons water per acre.

For Air Application: Use recommended amounts in 10-20 gallons water per acre.

DIRECTIONS

ARTICHOKES: Plume Moth, Aphids—4 lbs. per acre in water to cover. Do not apply within 7 days of harvest.

BROCCOLI, CABBAGE, CAULIFLOWER: Plume Moth, Aphids, Armyworm Larvae (Up to 3rd Instar)—4 to 6 lbs. per acre in water to cover. Do not apply within 21 days of harvest.

ACCEPTED

Nov. 5, 1971

239-2074



POISON



ANTIDOTE: If Swallowed. Give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Have victim lie down and keep quiet. **CALL A PHYSICIAN IMMEDIATELY!** Note to Physicians: Atropine is Antidotal. 2-PAM is also antidotal and may be administered in conjunction with atropine. If On Skin. In case of contact, remove contaminated clothing and immediately wash skin with soap and water.

DANGER:

KEEP OUT OF REACH OF CHILDREN
DO NOT TAKE INTERNALLY
DO NOT INHALE
DO NOT GET ON SKIN

DANGER:

Poisonous If Swallowed, Inhaled, or Absorbed Through Skin. Wear a mask or respirator of a type passed by the U. S. Bureau of Mines for methyl parathion protection. Rapidly Absorbed Through Skin! Do not get in eyes, on skin or on clothing. Wear natural rubber gloves, protective clothing and goggles. In case of contact, wash immediately with soap and water. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use. Keep all unprotected persons out of operating areas or vicinity where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticides and volatile residues have dissipated. Do not store or transport near feed or food. Do not use on sorghum (milo). Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

Toxic to wildlife. Birds feeding on treated areas may be killed. Do not apply when weather conditions favor drift

BRUSSELS SPROUTS: Armyworm Larvae (Up to 3rd Instar)—4 lbs. per acre in water to cover. Do not apply within 21 days of harvest.

CELERY, PEAS, PEPPERS, TOMATOES: Aphids, Loopers, Armyworm Larvae (Up to 3rd Instar)—4 lbs. per acre in water to cover. Do not apply within 15 days of harvest.

LETTUCE, SPINACH: Aphids—4 lbs. per acre in water to cover. Do not apply within 21 days of harvest.

CONDITIONS OF SALE: 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed hereon. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material. 2. Critical and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control and damage caused by drift to other plants or crops. Such risks occur even though the product is reasonably fit for the uses stated hereon and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability (except those assumed by Chevron under 1 above) resulting from handling, storage, and use of this material.

from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Do not contaminate any body of water, woodlands, or other non-crop areas. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.