

KELTHANE-Sulfur 4-50 Dust

(MITICIDE-FUNGICIDE)

Active Ingredients
1,1-bis p-chlorophenyl-2,2,2-trichloroethanol
Sulfur
Inert Ingredients

By Wt.
4%
50%
46%

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

THIS PRODUCT IS A DUST. DO NOT SPRAY OR SPOT TREAT. DO NOT USE ON VEGETABLES OR FRUIT. DO NOT USE ON CLOTHES, HAIR, ETC.

DO NOT USE ON CLOTHES, HAIR, ETC.

DIRECTIONS

ALFALFA: Seed, Cowpea, Southern Mite, Strawberry: Atlantic
Mite, Turnip Mite, Pacific Mite

APPLES: Spider Mites—60 to 70 lbs. per acre. Apply at petal fall to second cover, or whenever mite population requires. Repeat application if necessary. Wait 3 to 12 days between applications. Apply spray when 10% to 15% of leaves are infested with the first few mites.

BEANS: Spider Mites—30 to 37 lbs. per acre by ground application. Do not feed treated vines to cattle or horses. Do not apply within 7 days of harvest.

FIGS: Red Spider Mites—50 lbs. per acre. Apply at petal fall to first cover, or whenever mite population requires. Repeat application every 7 to 10 days. Do not apply within 7 days of harvest.

NET WEIGHT 50 LBS.

Chevron Chemical Company



ORTHO

ACCEPTED

JUN 20, 73

REVIEW THE PESTICIDE USE REQUIREMENTS FOR MITICIDE AND FUNGICIDE USE. USE THIS PRODUCT FOR ECONOMY AND EFFICIENCY. DO NOT USE ON CLOTHES, HAIR, ETC. TO ATTACH A LIQUIDANT.

GRAPES: Powdery Mildew, Spider Mites—30 to 37 lbs. per acre by ground application. Repeat application every 7 to 10 days. Do not apply within 7 days of harvest.

HOPS: Two-spotted and Pacific Mite—50 lbs. per acre. Apply at petal fall to first cover, or whenever mite population requires. Do not apply within 7 days of harvest.

STRAWBERRIES: Red Spider Mites, Powdery Mildew—50 lbs. per acre. Apply at petal fall to first cover, or whenever mite population requires. Do not apply within 7 days of harvest.

CONDITIONS OF SALE: This product is sold subject to the conditions of sale set forth in the accompanying Sales Agreement.



READ THE LABEL

KEEP OUT OF THE REACH OF CHILDREN

CAUTION:

Do not apply to blossoms from the time of bloom until 14 days after bloom, or until 14 days before other planting. Do not apply to blossoms on the same plant during the same period.