

DIBROM[®] Sulfur 4-10 Dust [239-2145]



ORTHO ACCEPTED

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTRATION
E.O. UNDER NO. 11735
RECORDED, DOMESTIC, 5/12/68

(INSECTICIDE-FUNGICIDE)

Active Ingredients	By Wt.
*Naled	4%
Sulfur	10%
Inert Ingredients	86%
*1,2-dibromo-2,2-dichloroethyl dimethyl phosphate U. S. Pat. No. 2,971,882	

Use within _____ days of manufacture only, as this product decomposes.

Date of Manufacture _____

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER.

STORE IN COOL, DRY PLACE,
OUT OF DIRECT SUNLIGHT

BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE.

DIRECTIONS

Use 40 to 50 pounds per acre unless otherwise specified. This product is formulated for application by ground or air equipment. Begin applications at first sign of insects or disease. Repeat as necessary. Apply for contact and thorough coverage to obtain control of insects.

BEANS: Aphids, Mites, Lygus Bugs, Thrips, Leafhoppers, Leafminers, Loopers—50 lbs. per acre by ground or air. Do not apply within 4 days of harvest.

NET WEIGHT 50 LBS.

Chevron Chemical Company

Ortho Division, San Francisco, Calif. 94119

Richmond, California Fresno, California Honolulu, Hawaii
Des Moines, Iowa Perth Amboy, New Jersey Orlando, Florida
Form 6856 I Product 3968 Made in U.S.A.

EPA Reg. No. 239-2145-AA

BROCCOLI, CABBAGE, CAULIFLOWER, COLLARDS, BRUSSELS SPROUTS, KALE: Cabbage Looper, Imported Cabbageworm, Diamond-back Caterpillar, Aphids. May be used 1 day before harvest on Broccoli, Cabbage, Cauliflower and Brussels Sprouts. Do not apply to Collards and Kale within 4 days of harvest.

CITRUS (Oranges, Lemons, Grapefruit): Aphids, Red Spider Mites, Citrus Cutworms, Thrips—Do not apply within 7 days of harvest. California and Arizona—do not apply within 1 day of harvest. Do not apply in citrus groves adjacent to estuaries, as shrimp and crab may be killed at application rates recommended on this label.

COTTON: Lygus, Cotton Leaf Perforator, Armyworms, Looper, Fleahoppers—35 lbs. per acre. Apply at 7 to 10 days before first picking. Do not apply within 4 days of harvest.

GRAPES: Leafhoppers, Powdery Mildew—40 to 50 lbs. per acre. Thrips—20 to 25 lbs. per acre. Do not apply within 4 days of harvest.


LETTUCE, SPINACH, CELERY, CHARD, TURNIP GREENS: Loopers, Aphids—37 lbs. per acre. May be used 1 day before harvest on Lettuce, Spinach, Turnip Greens and Chard. Do not apply to Celery within 4 days of harvest.

PEACHES: Western States Only—Fruit Flies (*Drosophila*), Red Spider Mites, Peach Twig Borer, Lygus Bugs, Thrips in Calif. only, Powdery Mildew—50 lbs. per acre. Do not apply within 4 days of harvest.

STRAWBERRIES: Spider Mites, Aphids, Powdery Mildew. Do not apply within 4 days of harvest.

TOMATOES (Field): Two-Spotted Mites, Aphids, Leaf Miners, Tomato Fruitworms, Hornworms, Tomato Russet Mites—25 lbs. per acre. Make first application at 5 to 7 days before first picking and continue at 5 to 7-day intervals until harvest is complete. Do not apply within 1 day of harvest. Because of the sulfur content of this dust, first obtain permission from your canner or processor before using it on tomatoes intended for sale to such canner or processor.

CONDITIONS OF SALE: 1. Chevron Chemical Company (Chevron) warrants that this material conforms to the chemical description on the label and is reasonably fit for use as directed herein. Chevron neither makes, nor authorizes any agent or representative to make, any other warranty of FITNESS or of MERCHANTABILITY, guarantee or representation, express or implied, concerning this material. 2. Critical and unforeseeable factors beyond Chevron's control prevent it from eliminating all risks in connection with the use of chemicals. Such risks include, but are not limited to, damage to plants and crops to which the material is applied, lack of complete control, and damage caused by drift to other plants and crops. Such risks occur even though the product is reasonably fit for the uses stated herein and even though label directions are followed. Buyer and user acknowledge and assume all risks and liability except those caused by fraud or under applicable laws that may be applicable to the application of this material.



READ THE LABEL

KEEP OUT OF REACH OF CHILDREN

CAUTION: Harmful if swallowed. May cause irritation of eyes, nose and throat. Avoid breathing dust. Avoid contact with skin and eyes. Wash thoroughly after using. Do not store or transport near feed or food. **Note to Physician:** Atropine is antidotal. PAM is also an antidote and may be administered in conjunction with atropine. Do not use on cherries as injury may result. May corrode metal dust equipment, clean equipment after use. Some fruits and other plants are susceptible to injury from sulfur under certain climatic conditions. The user is advised not to use sulfur on any crop unless local use has proved that sulfur does not damage crops in that locality. **Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.**

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

This product is highly toxic to bees exposed to direct treatment or drift on crops. Protective information may be obtained from your cooperative Agricultural Extension Station.