



SOD WEBWORM spray

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

ORTHO Sod Webworm Spray containing Lindane is designed for the control of Sod Webworms and other insects that infest lawns. It may be applied effectively with the ORTHO Lawn Sprayer.

Sod Webworms are light brown caterpillars, $\frac{1}{4}$ to $\frac{3}{4}$ inch long, that hide in the grass during the day and feed on the grass blades at night. Adult Webworms, usually referred to as Lawn Moths (about one inch in length with long snout), emerge from the grass at dusk. They fly over the lawn laying eggs that later hatch into destructive caterpillars.

Sod Webworm damage appears first as small brown spots which may rapidly develop into large brown areas of severely damaged or destroyed grass.

DIRECTIONS

Treat lawns at first sign of insect infestation. Repeat as necessary. Do not store diluted spray. Make new dilution for each use.

PREVENTION: To prevent damage from occurring, spray the lawn as directed monthly throughout the summer starting in late June.

CONTROL: If Lawn Moths (Sod Webworm Adults) are noticed in large numbers at dusk or at first sign of damage to lawn, treat as directed in following table. Repeat at monthly intervals throughout the summer.

PESTS	AMOUNT OF PRODUCT TO USE	COMMENTS
Sod Webworm	1 Cupful (8 fluid ozs.) in 15 gals. of water to treat 1,000 sq. ft.	If lawn is dry, water well before treatment. Apply in early evening when caterpillars are feeding. Heavy watering or irrigation should be withheld for 2 to 3 days after treatment.
Grasshoppers, Army worms, Cutworms, Fleas, Earwigs	1 Cupful (8 fluid ozs.) in 15 gals. of water to treat 1,000 sq. ft.	If lawn is dry, water well before treatment. Heavy watering or irrigation should be withheld for 2 to 3 days after treatment.
Ticks	5 Tbsp. (15 fluid ozs.) in 15 gals. of water to treat 1,000 sq. ft.	
White Grub	1 Cupful (8 fluid ozs.) in 15 gals. of water to treat 1,000 sq. ft.	Immediately after treatment, water lawn thoroughly to carry insecticide down into the root zone where insects are feeding.

HOW TO APPLY

Place required amount of product for 1,000 square feet (20' x 50') in the ORTHO Lawn Sprayer jar. (See table above for details). Fill remainder of jar with water. Apply evenly to 1,000 square feet of lawn.

CAUTION: Harmful if swallowed. Induce vomiting if swallowed. Avoid breathing spray mist. May be absorbed through skin. Avoid skin contact. In case of skin contact, wash immediately with soap and water. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Keep children and pets off treated areas until this material is washed into the soil and grass is dry. **DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.** When container is empty, immediately wash thoroughly and destroy. Never re-use.

This product is toxic to fish and wildlife.

Do not contaminate ornamental fish ponds. Note: Birds feeding in treated areas may be harmed.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.



CHEVRON CHEMICAL COMPANY

ORTHO DIVISION San Francisco Calif. 94109

RICHMOND, CALIF. Atlanta, Georgia Des Plaines, Illinois
Form 7630 A USDA Reg. No. 239-2266 Product 3914 Made in U.S.A.

SOD WEBWING
spray

USE STRICTLY IN ACCORDANCE WITH
LABEL DIRECTIONS.

DO NOT USE IN OR NEAR
FOOD OR DRINK CONTAINERS.

Apply to the body of the insect
or to the surface of the material
on which it is present.
Waterproofing is not necessary.
The day after treatment,
wash the treated surface with
water to remove the residue.

STAY AWAY FROM CHILDREN.

CAUTIONS:
Do not use in or near
food or drink containers.

Do not use in or near
children's play areas.
Do not use in or near
pets.

Ortho SOD Webwing Spray


HOW TO APPLY

Apply to the body of the insect
or to the surface of the material
on which it is present.
Waterproofing is not necessary.
The day after treatment,
wash the treated surface with
water to remove the residue.


CAUTION

Do not use in or near
food or drink containers.
Do not use in or near
children's play areas.
Do not use in or near
pets.

THE CHEVRON CHEMICAL COMPANY



SOD WEBWING
spray



ALSO KILLS:

Fleas, Ticks



SOD WEBWORM Spray

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

ORTHO Sod Webworm Spray containing Lindane is designed for the control of Sod Webworms and other insects that infest lawns. It may be applied effectively with the ORTHO Lawn Sprayer.

Sod Webworms are light brown caterpillars, 1/4 to 3/4 inch long, that hide in the grass during the day and feed on the grass blades at night. Adult Webworms, usually referred to as Lawn Moths (about one inch in length with long snout), emerge from the grass at dusk. They fly over the lawn laying eggs that later hatch into destructive caterpillars.

Sod Webworm damage appears first as small brown spots which may rapidly develop into large brown areas of severely damaged or destroyed grass.

DIRECTIONS

Treat lawns at first sign of insect infestation. Repeat as necessary. Do not store diluted spray. Make new dilution for each use.

PREVENTION: To prevent damage from occurring, spray the lawn as directed monthly throughout the summer starting in late June.

CONTROL: If Lawn Moths (Sod Webworm Adults) are noticed in large numbers at dusk or at first sign of damage to lawn, treat as directed in following table. Repeat at monthly intervals throughout the summer.

RECEIVED
Apr 6, 1970

PESTS	AMOUNT OF PRODUCT TO USE	COMMENTS
Sod Webworms 239-2260	1 Cupful (8 fluid ozs.) in 15 gals. of water to treat 1,000 sq. ft.	If lawn is dry, water well before treatment. Apply in early evening when caterpillars are feeding. Heavy watering or irrigation should be withheld for 2 to 3 days after treatment.
Grasshoppers, Armyworms, Cutworms, Fleas, Earwigs	1 Cupful (8 fluid ozs.) in 15 gals. of water to treat 1,000 sq. ft.	If lawn is dry, water well before treatment. Heavy watering or irrigation should be withheld for 2 to 3 days after treatment.
Ticks	3 Tbsp. (1.5 fluid ozs.) in 15 gals. of water to treat 1,000 sq. ft.	
White Grubs	1 Cupful (8 fluid ozs.) in 15 gals. of water to treat 1,000 sq. ft.	Immediately after treatment, water lawn thoroughly to carry insecticide down into the root zone where insects are feeding.

HOW

Place required amount of product in the ORTHO Lawn Sprayer jar. Fill remainder of jar with water. Apply

CAUTION: Harmful if swallowed or inhaled. Avoid breathing spray mist. May be absorbed through skin contact. In case of skin contact, wash with soap and water. Avoid contamination of feed and pets or humans. Keep children away from treated areas. If material is washed into the soil

SPILL OR STORE NEAR HEAT

empty, immediately wash thoroughly. This product is toxic to fish and aquatic life.

Do not contaminate ornamentals. Treated areas may be harmed.

NOTICE: Buyer assumes all responsibility in accordance with directions.



CHEVRON
ORTHODIVISION

RICHMOND, CALIF.
Form 7631 A

Attn:
USDA Reg. No. 239-22